

# Schatten-class perturbations of Toeplitz operators

Dominik Schillo

Saarland University

Multivariable Operator Theory, 2017

Joint work with  
Michael Didas and Jörg Eschmeier  
(both Saarland University)

## Definition

Let  $\mathbb{T}$  be the unit circle in  $\mathbb{C}$  with the canonical probability measure  $m$ . The *Hardy space with respect to  $m$*  will be denoted by

$$H^2(m) = \left\{ f \in L^2(m) ; \hat{f}(n) = 0 \text{ for all } n < 0 \right\} \subset L^2(m).$$

## Definition

Let  $\mathbb{T}$  be the unit circle in  $\mathbb{C}$  with the canonical probability measure  $m$ . The *Hardy space with respect to  $m$*  will be denoted by

$$H^2(m) = \left\{ f \in L^2(m) ; \hat{f}(n) = 0 \text{ for all } n < 0 \right\} \subset L^2(m).$$

Define

$$H^\infty(m) = L^\infty(m) \cap H^2(m) \subset L^\infty(m)$$

## Definition

Let  $\mathbb{T}$  be the unit circle in  $\mathbb{C}$  with the canonical probability measure  $m$ . The *Hardy space with respect to  $m$*  will be denoted by

$$H^2(m) = \left\{ f \in L^2(m) ; \hat{f}(n) = 0 \text{ for all } n < 0 \right\} \subset L^2(m).$$

Define

$$H^\infty(m) = L^\infty(m) \cap H^2(m) \subset L^\infty(m)$$

and we call

$$I_m = \{ f \in H^\infty(m) ; |f| = 1 \text{ } m\text{-a.e.} \}$$

the set of *inner functions with respect to  $m$* .

## Definition

Let  $f \in L^\infty(m)$ . We call

$$T_f: H^2(m) \rightarrow H^2(m), \quad g \mapsto P_{H^2(m)}(fg),$$

where  $P_{H^2(m)}: L^2(m) \rightarrow H^2(m)$  is the orthogonal projection onto  $H^2(m)$ , the *Toeplitz operator with symbol  $f$* .

## Theorem (Brown-Halmos condition I, 1964)

An operator  $X \in B(H^2(m))$  is a Toeplitz operator (i.e. there exists a  $f \in L^\infty(m)$  such that  $X = T_f$ ) if and only if

$$T_z^* X T_z - X = 0,$$

where  $z \in L^\infty(m)$  is the identity map.

## Theorem (Brown-Halmos condition I, 1964)

An operator  $X \in B(H^2(m))$  is a Toeplitz operator (i.e. there exists a  $f \in L^\infty(m)$  such that  $X = T_f$ ) if and only if

$$T_z^* X T_z - X = 0,$$

where  $z \in L^\infty(m)$  is the identity map.

## Theorem (Brown-Halmos condition II)

An operator  $X \in B(H^2(m))$  is a Toeplitz operator if and only if

$$T_u^* X T_u - X = 0$$

for all  $u \in I_m$ .

## Theorem (Gu, 2004)

*An operator  $X \in B(H^2(m))$  is a finite rank Toeplitz perturbation (i.e. there exist a  $f \in L^\infty(m)$  and  $F \in F(H^2(m))$ ) such that  $X = T_f + F$ ) if and only if*

$$T_u^* X T_u - X \in F(H^2(m))$$

*for all  $u \in I_m$ .*



## Theorem (Xia, 2009)

*An operator  $X \in B(H^2(m))$  is a compact Toeplitz perturbation (i.e. there exist a  $f \in L^\infty(m)$  and  $K \in K(H^2(m))$  such that  $X = T_f + K$ ) if and only if*

$$T_u^* X T_u - X \in K(H^2(m))$$

*for all  $u \in I_m$ .*

## Definition

Let  $p \in [1, \infty)$  and let  $H$  be a Hilbert space. An operator  $S \in B(H)$  is a *Schatten- $p$ -class* operator if

$$\|S\|_p^p = \operatorname{tr}(|S|^p) < \infty.$$

## Definition

Let  $p \in [1, \infty)$  and let  $H$  be a Hilbert space. An operator  $S \in B(H)$  is a *Schatten- $p$ -class operator* if

$$\|S\|_p^p = \operatorname{tr}(|S|^p) < \infty.$$

Furthermore, we set

$$\mathcal{S}_p(H) = \left\{ S \in B(H) ; \|S\|_p < \infty \right\}$$

equipped with  $\|\cdot\|_p$

## Definition

Let  $p \in [1, \infty)$  and let  $H$  be a Hilbert space. An operator  $S \in B(H)$  is a *Schatten- $p$ -class operator* if

$$\|S\|_p^p = \operatorname{tr}(|S|^p) < \infty.$$

Furthermore, we set

$$\mathcal{S}_p(H) = \left\{ S \in B(H) ; \|S\|_p < \infty \right\}$$

equipped with  $\|\cdot\|_p$  and

$$\mathcal{S}_0(H) = F(H) \quad \text{as well as} \quad \mathcal{S}_\infty(H) = K(H)$$

both equipped with the operator norm.

Let  $\mathbb{T} \subset \mathbb{C}$  be the unit circle with the canonical probability measure  $m \in M_1^+(\mathbb{T})$ .

Theorem (Gu, Xia, Didas-Eschmeier-S.)

Let  $p \in \{0\} \cup [1, \infty]$ . An operator  $X \in B(H^2(m))$  is a  $\mathcal{S}_p(H^2(m))$  Toeplitz perturbation (i.e. there exist a  $f \in L^\infty(m)$  and  $S \in \mathcal{S}_p(H^2(m))$  such that  $X = T_f + S$ ) if and only if

$$T_u^* X T_u - X \in \mathcal{S}_p(H^2(m))$$

for all  $u \in I_m$ .

Let  $\mathbb{T} \subset \mathbb{C}$  be the unit circle with the canonical probability measure  $m \in M_1^+(\mathbb{T})$ .

Theorem (Gu, Xia, Didas-Eschmeier-S.)

Let  $p \in \{0\} \cup [1, \infty]$ . An operator  $X \in B(H^2(m))$  is a  $S_p(H^2(m))$  Toeplitz perturbation (i.e. there exist a  $f \in L^\infty(m)$  and  $S \in \mathcal{S}_p(H^2(m))$  such that  $X = T_f + S$ ) if and only if

$$T_u^* X T_u - X \in \mathcal{S}_p(H^2(m))$$

for all  $u \in I_m$ .

## Definition

Let  $\mathbb{D}$  be the unit disc in  $\mathbb{C}$  and  $\mathcal{O}(\mathbb{D})$  be the set of all scalar-valued analytic functions on  $\mathbb{D}$ .

## Definition

Let  $\mathbb{D}$  be the unit disc in  $\mathbb{C}$  and  $\mathcal{O}(\mathbb{D})$  be the set of all scalar-valued analytic functions on  $\mathbb{D}$ . We call

$$A(\mathbb{D}) = \{f \in C(\overline{\mathbb{D}}) ; f|_{\mathbb{D}} \in \mathcal{O}(\mathbb{D})\}$$

the *disc algebra*



## Definition

Let  $\mathbb{D}$  be the unit disc in  $\mathbb{C}$  and  $\mathcal{O}(\mathbb{D})$  be the set of all scalar-valued analytic functions on  $\mathbb{D}$ . We call

$$A(\mathbb{D}) = \{f \in C(\overline{\mathbb{D}}) ; f|_{\mathbb{D}} \in \mathcal{O}(\mathbb{D})\}$$

the *disc algebra* and denote by  $\partial_{A(\mathbb{D})}$  the *Shilov boundary* of  $A(\mathbb{D})$  (i.e. the smallest closed subset of  $\overline{\mathbb{D}}$  such that

$$\sup_{z \in \overline{\mathbb{D}}} |f(z)| = \sup_{z \in \partial_{A(\mathbb{D})}} |f(z)| \quad \text{for all } f \in A(\mathbb{D}).$$

## Definition

Let  $\mathbb{D}$  be the unit disc in  $\mathbb{C}$  and  $\mathcal{O}(\mathbb{D})$  be the set of all scalar-valued analytic functions on  $\mathbb{D}$ . We call

$$A(\mathbb{D}) = \{f \in C(\overline{\mathbb{D}}) ; f|_{\mathbb{D}} \in \mathcal{O}(\mathbb{D})\}$$

the *disc algebra* and denote by  $\partial_{A(\mathbb{D})}$  the *Shilov boundary* of  $A(\mathbb{D})$  (i.e. the smallest closed subset of  $\overline{\mathbb{D}}$  such that

$$\sup_{z \in \overline{\mathbb{D}}} |f(z)| = \sup_{z \in \partial_{A(\mathbb{D})}} |f(z)| \quad \text{for all } f \in A(\mathbb{D}).$$

## Proposition

We have

$$\partial_{A(\mathbb{D})} = \partial\mathbb{D} = \mathbb{T}.$$

## Definition

Let  $D \subset \mathbb{C}^n$  be a bounded domain. We denote by

- (i)  $A(D) = \{f \in C(\overline{D}) ; f|_D \in \mathcal{O}(D)\} \subset C(\overline{D})$  the *domain algebra* of  $D$ ,

## Definition

Let  $D \subset \mathbb{C}^n$  be a bounded domain. We denote by

- (i)  $A(D) = \{f \in C(\overline{D}) ; f|_D \in \mathcal{O}(D)\} \subset C(\overline{D})$  the *domain algebra* of  $D$ ,
- (ii)  $S = \partial_{A(D)}$  the *Shilov boundary* of  $A(D)$  (i.e. the smallest closed subset of  $\overline{D}$  such that

$$\sup_{z \in \overline{D}} |f(z)| = \sup_{z \in S} |f(z)|$$

for all  $f \in A(D)$ ).

Let  $D \subset \mathbb{C}^n$  be a strictly pseudoconvex or a bounded symmetric and circled domain. We denote by  $\mu \in M_1^+(S)$  the canonical probability measure on  $S$ .

Let  $D \subset \mathbb{C}^n$  be a strictly pseudoconvex or a bounded symmetric and circled domain. We denote by  $\mu \in M_1^+(S)$  the canonical probability measure on  $S$ .

## Examples

(i)  $D = \mathbb{B}_n$ :  $S = \partial\mathbb{B}_n$  and  $\mu = \sigma$ .

(ii)  $D = \mathbb{D}^n$ :  $S = \mathbb{T}^n$  and  $\mu = \otimes_n m$ .

We have

$$H^2(m) = \overline{A(\mathbb{D})|_{\mathbb{T}}^{\tau_{\|\cdot\|_{L^2(m)}}}} \quad \text{and} \quad H^\infty(m) = \overline{A(\mathbb{D})|_{\mathbb{T}}^{\tau_{w^*}}}.$$

We have

$$H^2(m) = \overline{A(\mathbb{D})|_{\mathbb{T}}}^{\tau_{\|\cdot\|}} L^2(m) \quad \text{and} \quad H^\infty(m) = \overline{A(\mathbb{D})|_{\mathbb{T}}}^{\tau_{w^*}}.$$

### Definition

We define

$$H^2(\mu) = \overline{A(D)|_S}^{\tau_{\|\cdot\|}} L^2(\mu) \subset L^2(\mu)$$

and

$$H^\infty(\mu) = \overline{A(D)|_S}^{\tau_{w^*}} \subset L^\infty(\mu).$$



We have

$$H^2(m) = \overline{A(\mathbb{D})|_{\mathbb{T}}}_{\tau_{\|\cdot\|_{L^2(m)}}} \quad \text{and} \quad H^\infty(m) = \overline{A(\mathbb{D})|_{\mathbb{T}}}_{\tau_{w^*}}.$$

## Definition

We define

$$H^2(\mu) = \overline{A(D)|_S}_{\tau_{\|\cdot\|_{L^2(\mu)}}} \subset L^2(\mu)$$

and

$$H^\infty(\mu) = \overline{A(D)|_S}_{\tau_{w^*}} \subset L^\infty(\mu).$$

Furthermore, we denote by

$$I_\mu = \{f \in H^\infty(\mu) ; |f| = 1 \text{ } \mu\text{-a.e.}\}$$

the set of inner functions with respect to  $\mu$ .

## Definition

Let  $f \in L^\infty(\mu)$ . We call

$$T_f: H^2(\mu) \rightarrow H^2(\mu), \quad g \mapsto P_{H^2(\mu)}(fg),$$

where  $P_{H^2(\mu)}: L^2(\mu) \rightarrow H^2(\mu)$  is the orthogonal projection onto  $H^2(\mu)$ , the *Toeplitz operator with symbol  $f$* .

## Definition

Let  $f \in L^\infty(\mu)$ . We call

$$T_f: H^2(\mu) \rightarrow H^2(\mu), \quad g \mapsto P_{H^2(\mu)}(fg),$$

where  $P_{H^2(\mu)}: L^2(\mu) \rightarrow H^2(\mu)$  is the orthogonal projection onto  $H^2(\mu)$ , the *Toeplitz operator with symbol  $f$* .

## Theorem (Didas-Eschmeier-Everard, 2011)

*An operator  $X \in B(H^2(\mu))$  is a Toeplitz operator if and only if*

$$T_u^* X T_u - X = 0$$

*for all  $u \in I_\mu$ .*

## Definition

We denote by  $H^\infty(\mathbb{D}) \subset \mathcal{O}(\mathbb{D})$  the set of all bounded analytic functions on  $\mathbb{D}$ .

## Definition

We denote by  $H^\infty(\mathbb{D}) \subset \mathcal{O}(\mathbb{D})$  the set of all bounded analytic functions on  $\mathbb{D}$ .

## Theorem

*The map*

$$r_m: H^\infty(\mathbb{D}) \rightarrow H^\infty(m), \theta \mapsto \tau_{w^*}\text{-}\lim_{r \rightarrow 1} [\theta(r \cdot)] =: \theta^*$$

*is an isometric algebra isomorphism and weak\* homeomorphism with  $r_m(\theta|_{\mathbb{D}}) = [\theta|_{\mathbb{T}}]$  for all  $\theta \in A(\mathbb{D})$ .*

## Definition

We denote by  $H^\infty(D) \subset \mathcal{O}(D)$  the set of all bounded analytic functions on  $D$ .

## Definition

We denote by  $H^\infty(D) \subset \mathcal{O}(D)$  the set of all bounded analytic functions on  $D$ .

## Theorem

*There exists a map*

$$r_\mu: H^\infty(D) \rightarrow H^\infty(\mu), \theta \mapsto r_\mu(\theta) =: \theta^*,$$

*which is an isometric algebra isomorphism and weak\* homeomorphism with  $r_\mu(\theta|_D) = [\theta|_S]$  for all  $\theta \in A(D)$ .*

Let  $D \subset \mathbb{C}^n$  be a strictly pseudoconvex or bounded symmetric and circled domain and  $\mu \in M_1^+(S)$  be the probability measure on the Shilov boundary  $S = \partial_{A(D)}$  obtained before.

### Theorem (Didas-Eschmeier-S., 2017)

*Let  $p \in [1, \infty)$ . An operator  $X \in B(H^2(\mu))$  is a  $\mathcal{S}_p(H^2(\mu))$  Toeplitz perturbation (i.e. there exist a  $f \in L^\infty(\mu)$  and  $S \in \mathcal{S}_p(H^2(\mu))$  such that  $X = T_f + S$ ) if and only if*

$$T_u^* X T_u - X \in \mathcal{S}_p(H^2(\mu))$$

*for all  $u \in I_\mu$ .*



## Proposition

Let  $(\alpha_k)_{k \in \mathbb{N}}$  be a sequence in  $H^\infty(\mu)$  with

$$\tau_{w^*}\text{-}\lim_{k \rightarrow \infty} \alpha_k = \alpha \in [0, 1)$$

## Proposition

Let  $(\alpha_k)_{k \in \mathbb{N}}$  be a sequence in  $H^\infty(\mu)$  with

$$\tau_{w^*}\text{-}\lim_{k \rightarrow \infty} \alpha_k = \alpha \in [0, 1)$$

and  $X \in B(H^2(\mu))$  an operator such that

$$Y = \tau_{\text{WOT}}\text{-}\lim_{k \rightarrow \infty} T_{\alpha_k}^* X T_{\alpha_k} \in B(H^2(\mu))$$

exists.

## Proposition

Let  $(\alpha_k)_{k \in \mathbb{N}}$  be a sequence in  $H^\infty(\mu)$  with

$$\tau_{w^*}\text{-}\lim_{k \rightarrow \infty} \alpha_k = \alpha \in [0, 1)$$

and  $X \in B(H^2(\mu))$  an operator such that

$$Y = \tau_{\text{WOT}}\text{-}\lim_{k \rightarrow \infty} T_{\alpha_k}^* X T_{\alpha_k} \in B(H^2(\mu))$$

exists. If  $T_u^* X T_u - X \in \mathcal{S}_\infty(H^2(\mu))$  for all  $u \in I_\mu$ , then there exists a function  $f \in L^\infty(\mu)$  such that

$$X = T_f + \frac{1}{1 - \alpha^2} (X - Y).$$

## Proposition

Let  $(\alpha_k)_{k \in \mathbb{N}}$  be a sequence in  $H^\infty(\mu)$  with

$$\tau_{W^*}\text{-}\lim_{k \rightarrow \infty} \alpha_k = \alpha \in [0, 1)$$

and  $X \in B(H^2(\mu))$  an operator such that

$$Y = \tau_{\text{WOT}}\text{-}\lim_{k \rightarrow \infty} T_{\alpha_k}^* X T_{\alpha_k} \in B(H^2(\mu))$$

exists. If  $T_u^* X T_u - X \in \mathcal{S}_\infty(H^2(\mu))$  for all  $u \in I_\mu$ , then there exists a function  $f \in L^\infty(\mu)$  such that

$$X = T_f + \frac{1}{1 - \alpha^2} (X - Y).$$

## Proposition (Hiai, 1997)

*The map*

$$\|\cdot\|_p : (B(H^2(\mu)), \tau_{\text{WOT}}) \rightarrow [0, \infty], S \mapsto \|S\|_p$$

*is lower semi-continuous.*

## Proposition (Hiai, 1997)

The map

$$\|\cdot\|_p : (B(H^2(\mu)), \tau_{\text{WOT}}) \rightarrow [0, \infty], S \mapsto \|S\|_p$$

is lower semi-continuous.

$$\left\| \tau_{\text{WOT}}\text{-}\lim_{k \rightarrow \infty} X - T_{\alpha_k}^* X T_{\alpha_k} \right\|_p \leq \liminf_{k \rightarrow \infty} \|X - T_{\alpha_k}^* X T_{\alpha_k}\|_p$$

We denote by

$$I_D = \{\theta \in H^\infty(D) ; \theta^* \in I_\mu\}$$

the set of inner functions with respect to  $D$  and  $\mu$ .

We denote by

$$I_D = \{\theta \in H^\infty(D) ; \theta^* \in I_\mu\}$$

the set of inner functions with respect to  $D$  and  $\mu$ .

### Proposition (Aleksandrov, 1984)

Let  $\alpha \in [0, 1)$ . Then there exists a sequence  $(\alpha_k)_{k \in \mathbb{N}}$  in  $I_D$  such that

$$\tau_{w^*}\text{-}\lim_{k \rightarrow \infty} \alpha_k^* = \alpha.$$



## Proposition (Xia, 2009)

Let  $p \in [1, \infty]$ . Suppose that  $X \in B(H^2(\mu))$  is an operator such that

$$T_u^* X T_u - X \in \mathcal{S}_p(H^2(\mu))$$

for all  $u \in I_\mu$ .

## Proposition (Xia, 2009)

Let  $p \in [1, \infty]$ . Suppose that  $X \in B(H^2(\mu))$  is an operator such that

$$T_u^* X T_u - X \in \mathcal{S}_p(H^2(\mu))$$

for all  $u \in I_\mu$ . Then, for all  $\varepsilon > 0$ , there is a  $\delta = \delta(\varepsilon) > 0$  such that

$$\|T_{\theta^*}^* X T_{\theta^*} - X\|_p \leq \varepsilon$$

for all  $\theta \in I_D$  with  $|\int_S 1 - \theta^* d\mu| \leq \delta$ .

## Proof.

There exists  $0 < \delta < 1$  such that  $\|T_{\theta^*}^* X T_{\theta^*} - X\|_p \leq 1$  for all  $\theta \in I_D$  with  $|\int_S 1 - \theta^* d\mu| \leq \delta$ . Set  $\alpha = 1 - \delta/2$ .

## Proof.

There exists  $0 < \delta < 1$  such that  $\|T_{\theta^*}^* X T_{\theta^*} - X\|_p \leq 1$  for all  $\theta \in I_D$  with  $|\int_S 1 - \theta^* d\mu| \leq \delta$ . Set  $\alpha = 1 - \delta/2$ .

$\implies$  There exists  $(\alpha_k)_{k \in \mathbb{N}}$  in  $I_D$  with  $\tau_{w^*}\text{-}\lim_{k \rightarrow \infty} \alpha_k^* = \alpha$ .

## Proof.

There exists  $0 < \delta < 1$  such that  $\|T_{\theta^*}^* X T_{\theta^*} - X\|_p \leq 1$  for all  $\theta \in I_D$  with  $|\int_S 1 - \theta^* d\mu| \leq \delta$ . Set  $\alpha = 1 - \delta/2$ .

$\implies$  There exists  $(\alpha_k)_{k \in \mathbb{N}}$  in  $I_D$  with  $\tau_{W^*}\text{-}\lim_{k \rightarrow \infty} \alpha_k^* = \alpha$ .

By passing to a subsequence we can achieve that

$|\int_D 1 - \alpha_k^* d\mu| \leq \delta$  for all  $k \in \mathbb{N}$  and that at the same time the limit

$$Y = \tau_{WOT}\text{-}\lim_{k \rightarrow \infty} T_{\alpha_k^*}^* X T_{\alpha_k^*} \in B(H^2(\mu))$$

exists.

## Proof.

There exists  $0 < \delta < 1$  such that  $\|T_{\theta^*}^* X T_{\theta^*} - X\|_p \leq 1$  for all  $\theta \in I_D$  with  $|\int_S 1 - \theta^* d\mu| \leq \delta$ . Set  $\alpha = 1 - \delta/2$ .

$\implies$  There exists  $(\alpha_k)_{k \in \mathbb{N}}$  in  $I_D$  with  $\tau_{W^*}\text{-}\lim_{k \rightarrow \infty} \alpha_k^* = \alpha$ .

By passing to a subsequence we can achieve that

$|\int_D 1 - \alpha_k^* d\mu| \leq \delta$  for all  $k \in \mathbb{N}$  and that at the same time the limit

$$Y = \tau_{WOT}\text{-}\lim_{k \rightarrow \infty} T_{\alpha_k^*}^* X T_{\alpha_k^*} \in B(H^2(\mu))$$

exists. Hence

$$\left\| \tau_{WOT}\text{-}\lim_{k \rightarrow \infty} X - T_{\alpha_k^*}^* X T_{\alpha_k^*} \right\|_p \leq \liminf_{k \rightarrow \infty} \left\| X - T_{\alpha_k^*}^* X T_{\alpha_k^*} \right\|_p \leq 1.$$

## Proof.

There exists  $0 < \delta < 1$  such that  $\|T_{\theta^*}^* X T_{\theta^*} - X\|_p \leq 1$  for all  $\theta \in I_D$  with  $|\int_S 1 - \theta^* d\mu| \leq \delta$ . Set  $\alpha = 1 - \delta/2$ .

$\implies$  There exists  $(\alpha_k)_{k \in \mathbb{N}}$  in  $I_D$  with  $\tau_{W^*}\text{-}\lim_{k \rightarrow \infty} \alpha_k^* = \alpha$ .

By passing to a subsequence we can achieve that

$|\int_D 1 - \alpha_k^* d\mu| \leq \delta$  for all  $k \in \mathbb{N}$  and that at the same time the limit

$$Y = \tau_{WOT}\text{-}\lim_{k \rightarrow \infty} T_{\alpha_k^*}^* X T_{\alpha_k^*} \in B(H^2(\mu))$$

exists. Hence

$$\left\| \tau_{WOT}\text{-}\lim_{k \rightarrow \infty} X - T_{\alpha_k^*}^* X T_{\alpha_k^*} \right\|_p \leq \liminf_{k \rightarrow \infty} \left\| X - T_{\alpha_k^*}^* X T_{\alpha_k^*} \right\|_p \leq 1.$$

$$\implies X - Y \in \mathcal{S}_p(H^2(\mu)).$$



## Remark

The following ingredients are essential for the proof.



## Remark

The following ingredients are essential for the proof.

- (i) The triple  $(A(D)|_S, S, \mu)$  is *regular* (in the sense of Aleksandrov).

## Remark

The following ingredients are essential for the proof.

- (i) The triple  $(A(D)|_S, S, \mu)$  is *regular* (in the sense of Aleksandrov).
- (ii) The measure  $\mu$  is a *faithful Henkin measure*.

## Remark

The following ingredients are essential for the proof.

- (i) The triple  $(A(D)|_S, S, \mu)$  is *regular* (in the sense of Aleksandrov).
- (ii) The measure  $\mu$  is a *faithful Henkin measure*.

## Question

What about  $p = 0, \infty$ ?

## Theorem

*An operator  $X \in B(H^2(m))$  is an analytic Toeplitz operator (i.e. there exists a  $f \in H^\infty(m)$  such that  $X = T_f$ ) if and only if*

$$[X, T_g] = XT_g - T_gX = 0.$$

*for all  $g \in H^\infty(m)$ .*

## Theorem

*An operator  $X \in B(H^2(m))$  is an analytic Toeplitz operator (i.e. there exists a  $f \in H^\infty(m)$  such that  $X = T_f$ ) if and only if*

$$[X, T_g] = XT_g - T_gX = 0.$$

*for all  $g \in H^\infty(m)$ .*

## Theorem (Davidson, 1977)

*An operator  $X \in B(H^2(m))$  has the form  $X = T_f + K$  with  $f \in H^\infty(m) + C(\mathbb{T})$  and  $K \in K(H^2(m))$  if and only if*

$$XT_g - T_gX \in K(H^2(m)).$$

*for all  $g \in H^\infty(m)$ .*

## Theorem (Hartman, 1958)

We have

$$H^\infty(m) + C(\mathbb{T}) = \{f \in L^\infty(m) ; H_f \text{ is compact}\}$$

with  $H_f = (1 - P_{H^2(m)})M_f|_{H^2(m)}$  for  $f \in L^\infty(m)$ .

## Theorem (Hartman, 1958)

We have

$$H^\infty(m) + C(\mathbb{T}) = \{f \in L^\infty(m) ; H_f \text{ is compact}\}$$

with  $H_f = (1 - P_{H^2(m)})M_f|_{H^2(m)}$  for  $f \in L^\infty(m)$ .

## Theorem (Davidson, Hartman)

An operator  $X \in B(H^2(m))$  has the form  $X = T_f + K$  with  $f \in L^\infty(m)$  such that  $H_f$  is compact and  $K \in K(H^2(m))$  if and only if

$$XT_g - T_gX \in K(H^2(m)).$$

for all  $g \in H^\infty(m)$ .

Let  $n \geq 1$  and

$$D = \mathbb{B}_n.$$



Let  $n \geq 1$  and

$$D = \mathbb{B}_n.$$

### Theorem (Ding-Sun, 1997)

*An operator  $X \in B(H^2(\mu))$  has the form  $X = T_f + K$  for some  $f \in L^\infty(\mu)$  such that  $H_f$  is compact and  $K \in K(H^2(\mu))$  if and only if*

$$XT_g - T_gX \in K(H^2(\mu))$$

*for all  $g \in H^\infty(\mu)$ .*

Let  $n \geq 1$  and

$$D = \mathbb{B}_n \quad \text{or} \quad D = \mathbb{D}^n.$$

Let  $n \geq 1$  and

$$D = \mathbb{B}_n \quad \text{or} \quad D = \mathbb{D}^n.$$

### Theorem (Guo-Wang, 2006)

*An operator  $X \in B(H^2(\mu))$  has the form  $X = T_f + F$  for some  $f \in L^\infty(\mu)$  such that  $H_f$  is of finite rank and  $F \in F(H^2(\mu))$  if and only if*

$$XT_g - T_gX \in F(H^2(\mu))$$

*for all  $g \in H^\infty(\mu)$ .*

Let  $n \geq 1$  and

$$D = \mathbb{B}_n \quad \text{or} \quad D = \mathbb{D}^n.$$

Theorem (Didas-Eschmeier-S., 2017)

*An operator  $X \in B(H^2(\mu))$  has the form  $X = T_f + S$  for some  $f \in L^\infty(\mu)$  such that  $H_f$  is in the Schatten- $p$ -class and  $S \in \mathcal{S}_p(H^2(\mu))$  if and only if*

$$XT_g - T_gX \in \mathcal{S}_p(H^2(\mu))$$

*for all  $g \in H^\infty(\mu)$ .*