

# Hyperbolic Geometry on Noncommutative Polyballs

GELU POPESCU

University of Texas at San Antonio

Technion-Israel Institute of Technology, June 2017

## Papers

- **G. Popescu**, Free Pluriharmonic Functions on Noncommutative Polyballs, *Analysis & PDE*, **9** (2016).
- **G. Popescu**, Hyperbolic Geometry on Noncommutative Polyballs, [submitted](#).

# Noncommutative polyballs

- $B(\mathcal{H})^{n_1} \times_c \cdots \times_c B(\mathcal{H})^{n_k}$  denotes the set of all tuples  $\mathbf{X} = (X_1, \dots, X_k)$  with the property that the entries of  $X_s := (X_{s,1}, \dots, X_{s,n_s})$  are commuting with the entries of  $X_t := (X_{t,1}, \dots, X_{t,n_t})$  for any  $s, t \in \{1, \dots, k\}$ ,  $s \neq t$ .

- The open *polyball* :

$$\mathbf{P}_n(\mathcal{H}) := [B(\mathcal{H})^{n_1}]_1 \times_c \cdots \times_c [B(\mathcal{H})^{n_k}]_1,$$

where  $[B(\mathcal{H})^{n_i}]_1$  is the open unit ball

$$\{(X_{i,1}, \dots, X_{i,n_i}) \in B(\mathcal{H})^{n_i} : \|X_{i,1}X_{i,1}^* + \cdots + X_{i,n_i}X_{i,n_i}^*\| < 1\}.$$

# Noncommutative regular polyballs

- The *regular polyball* on the Hilbert space  $\mathcal{H}$  is defined by

$$\mathbf{B}_n(\mathcal{H}) := \{\mathbf{X} \in \mathbf{P}_n(\mathcal{H}) : \Delta_{\mathbf{X}}(I) > 0\},$$

where the *defect mapping*  $\Delta_{\mathbf{X}} : B(\mathcal{H}) \rightarrow B(\mathcal{H})$  is given by

$$\Delta_{\mathbf{X}} := (id - \Phi_{X_1}) \circ \cdots \circ (id - \Phi_{X_k}),$$

and  $\Phi_{X_i} : B(\mathcal{H}) \rightarrow B(\mathcal{H})$  is the completely positive linear map defined by

$$\Phi_{X_i}(Y) := \sum_{j=1}^{n_i} X_{i,j} Y X_{i,j}^*, \quad Y \in B(\mathcal{H}).$$

- (*Abstract*) *regular polyball*  $\mathbf{B}_n := \coprod_{\mathcal{H}} \mathbf{B}_n(\mathcal{H})$ .

## Universal models

- Let  $H_{n_i}$  be an  $n_i$ -dimensional complex Hilbert space with orthonormal basis  $e_1^i, \dots, e_{n_i}^i$ . The **full Fock space** of  $H_{n_i}$  is defined by

$$F^2(H_{n_i}) := \mathbb{C}1 \oplus \bigoplus_{s \geq 1} H_{n_i}^{\otimes s}.$$

- Let  $\mathbb{F}_{n_i}^+$  be the unital free semigroup on  $n_i$  generators  $g_1^i, \dots, g_{n_i}^i$  and the identity  $g_0^i$ . Set  $e_\alpha^i := e_{j_1}^i \otimes \dots \otimes e_{j_p}^i$  if  $\alpha = g_{j_1}^i \dots g_{j_p}^i \in \mathbb{F}_{n_i}^+$  and  $e_{g_0^i}^i := 1 \in \mathbb{C}$ .
- For each  $i \in \{1, \dots, k\}$  and  $j \in \{1, \dots, n_i\}$ , the **left creation operator**  $S_{i,j}$  on  $F^2(H_{n_i})$  is defined by setting

$$S_{i,j} e_\alpha^i := e_j^i \otimes e_\alpha^i, \quad \alpha \in \mathbb{F}_{n_i}^+.$$

# Universal models

## Definition

The operator  $\mathbf{S}_{i,j}$  acting on  $F^2(H_{n_1}) \otimes \cdots \otimes F^2(H_{n_k})$  is defined by

$$\mathbf{S}_{i,j} := \underbrace{I \otimes \cdots \otimes I}_{i-1 \text{ times}} \otimes S_{i,j} \otimes \underbrace{I \otimes \cdots \otimes I}_{k-i \text{ times}}.$$

- Similarly, we define the *right creation operator*  $R_{i,j} : F^2(H_{n_i}) \rightarrow F^2(H_{n_i})$  by setting  $R_{i,j}e_\alpha^i := e_\alpha^i \otimes e_j^i$  and the corresponding  $\mathbf{R}_{i,j}$ .
- The *noncommutative polyball algebra*  $\mathcal{A}_n$  (resp  $\mathcal{R}_n$ ) is the norm closed non-selfadjoint algebra generated by  $\{\mathbf{S}_{i,j}\}$  (resp.  $\{\mathbf{R}_{i,j}\}$ ) and the identity.

## Universal models

- The  $k$ -tuple  $\mathbf{S} := (\mathbf{S}_1, \dots, \mathbf{S}_k)$ , where  $\mathbf{S}_i := (\mathbf{S}_{i,1}, \dots, \mathbf{S}_{i,n_i})$ , is a pure element in the regular polyball  $\mathbf{B}_n(\otimes_{i=1}^k F^2(H_{n_i}))^-$  and plays the role of *universal model* for the abstract regular polyball.
- Let  $\mathbf{X} = (X_1, \dots, X_k) \in \mathbf{B}_n(\mathcal{H})$  with  $X_i := (X_{i,1}, \dots, X_{i,n_i})$ .
- Set  $X_{i,\alpha_i} := X_{i,j_1} \cdots X_{i,j_p}$  if  $\alpha_i = g_{j_1}^i \cdots g_{j_p}^i \in \mathbb{F}_{n_i}^+$  and  $X_{i,g_0^i} := I$ .
- If  $\alpha := (\alpha_1, \dots, \alpha_k) \in \mathbb{F}_{n_1}^+ \times \cdots \times \mathbb{F}_{n_k}^+$ , denote  $\mathbf{X}_\alpha := \mathbf{X}_{1,\alpha_1} \cdots \mathbf{X}_{k,\alpha_k}$ .

## Main results on free pluriharmonic functions

- Introduce and characterize the class of  **$k$ -multi-Toeplitz operators** on  $F^2(H_{n_1}) \otimes \cdots \otimes F^2(H_{n_k})$ .
- Characterize the bounded free  **$k$ -pluriharmonic functions** and solve the Dirichlet extension problem on regular polyballs.
- Give necessary and sufficient conditions for a function to be the Poisson transform of a completely bounded (resp. completely positive) map on  $C^*(\mathbf{S})$ , the  $C^*$ -algebra generated by the universal model of the polyball.
- Obtain Herglotz-Riesz representation theorems for free holomorphic functions with positive real parts on regular polyballs.



# $k$ -multi-Toeplitz operators

- **Brown and Halmos** (Crelle, 1963) proved :

## Theorem

A bounded linear operator  $T$  on the Hardy space  $H^2(\mathbb{D})$  is a Toeplitz operator if and only if  $S^*TS = T$ , where  $S$  is the unilateral shift.

## Definition

A bounded linear operator  $T$  on  $F^2(H_{n_1}) \otimes \dots \otimes F^2(H_{n_k})$  is called  **$k$ -multi-Toeplitz operator** with respect to the right universal model  $\mathbf{R} = \{\mathbf{R}_{i,j}\}$  if, for each  $i \in \{1, \dots, k\}$ ,

$$\mathbf{R}_{i,s}^* T \mathbf{R}_{i,t} = \delta_{st} T, \quad s, t \in \{1, \dots, n_i\}.$$

## $k$ -multi-Toeplitz operators

- Each  $k$ -multi-Toeplitz operator  $T$  has a uniquely determined formal power series in several variables.
- One can recapture  $T$  from its “Fourier series”.
- We characterize the noncommutative formal power series which are Fourier series of  $k$ -multi-Toeplitz operators.

### Theorem

The set of all  **$k$ -multi-Toeplitz operators** on  $\bigotimes_{i=1}^k F^2(H_{n_i})$  coincides with

$$\mathcal{T}_n := \text{span}\{\mathcal{A}_n^* \mathcal{A}_n\}^{-\text{SOT}} = \text{span}\{\mathcal{A}_n^* \mathcal{A}_n\}^{-\text{WOT}},$$

where  $\mathcal{A}_n$  is the noncommutative polyball algebra.

# Noncommutative Berezin kernels

- If  $\mathbf{X} = \{X_{i,j}\} \in \mathbf{B}_n(\mathcal{H})^-$ , define the **noncommutative Berezin kernel**

$$\mathbf{K}_{\mathbf{X}} : \mathcal{H} \rightarrow \left(\otimes_{i=1}^k F^2(H_{n_i})\right) \otimes \overline{\Delta_{\mathbf{X}}(I)^{1/2}(\mathcal{H})}$$

by setting

$$\mathbf{K}_{\mathbf{X}} h := \sum_{\beta_i \in \mathbb{F}_{n_i}^+} e_{\beta_1}^1 \otimes \cdots \otimes e_{\beta_k}^k \otimes \Delta_{\mathbf{X}}(I)^{1/2} X_{1,\beta_1}^* \cdots X_{k,\beta_k}^* h,$$

where the defect operator is given by

$$\Delta_{\mathbf{X}}(I) := (id - \Phi_{X_1}) \circ \cdots \circ (id - \Phi_{X_k})(I).$$

# Noncommutative Berezin transforms

- The **Berezin transform** at  $\mathbf{X} \in \mathbf{B}_n(\mathcal{H})$  is the map  $\mathcal{B}_{\mathbf{X}} : B(\otimes_{i=1}^k F^2(H_{n_i})) \rightarrow B(\mathcal{H})$  defined by

$$\mathcal{B}_{\mathbf{X}}[g] := \mathbf{K}_{\mathbf{X}}^*(g \otimes I_{\mathcal{H}})\mathbf{K}_{\mathbf{X}}, \quad g \in B(\otimes_{i=1}^k F^2(H_{n_i})).$$

- If  $g \in C^*(\mathbf{S})$ , the  $C^*$ -algebra generated by  $\mathbf{S}_{i,1}, \dots, \mathbf{S}_{i,n_i}$ , we define the Berezin transform at  $\mathbf{X} \in \mathbf{B}_n(\mathcal{H})^-$  by

$$\mathcal{B}_{\mathbf{X}}[g] := \lim_{r \rightarrow 1} \mathbf{K}_{r\mathbf{X}}^*(g \otimes I_{\mathcal{H}})\mathbf{K}_{r\mathbf{X}},$$

where the limit is in the operator norm topology.

- $\mathcal{B}_{\mathbf{X}}$  is a unital completely positive linear map such that

$$\mathcal{B}_{\mathbf{X}}(\mathbf{S}_{\alpha}\mathbf{S}_{\beta}^*) = \mathbf{X}_{\alpha}\mathbf{X}_{\beta}^*, \quad \alpha, \beta \in \mathbb{F}_{n_1}^+ \times \dots \times \mathbb{F}_{n_k}^+,$$

where  $\mathbf{S}_{\alpha} := \mathbf{S}_{1,\alpha_1} \cdots \mathbf{S}_{k,\alpha_k}$  if  $\alpha := (\alpha_1, \dots, \alpha_k)$ .

# Free $k$ -pluriharmonic functions

## Definition

A function  $F$  is called **free  $k$ -pluriharmonic** on the polyball  $\mathbf{B}_n$  if it has the form

$$F(\mathbf{X}) = \sum_{m_1 \in \mathbb{Z}} \cdots \sum_{m_k \in \mathbb{Z}} \sum_{(\alpha, \beta) \in \Lambda} a_{\alpha, \beta} X_{1, \alpha_1} \cdots X_{k, \alpha_k} X_{1, \beta_1}^* \cdots X_{k, \beta_k}^*,$$

where  $(\alpha, \beta) \in \Lambda$  iff  $\alpha = (\alpha_1, \dots, \alpha_k)$  and  $\beta = (\beta_1, \dots, \beta_k)$ , with  $\alpha_i, \beta_i \in \mathbb{F}_{n_i}^+$ ,  $|\alpha_i| = m_i^-$ ,  $|\beta_i| = m_i^+$ , and the series converge in the operator norm topology for any  $\mathbf{X} = (X_1, \dots, X_k) \in \mathbf{B}_n(\mathcal{H})$  and any Hilbert space  $\mathcal{H}$ .

- $F$  is bounded if  $\|F\| := \sup_{\mathbf{X} \in \mathbf{B}_n(\mathcal{H})} \|F(\mathbf{X})\| < \infty$ .

## Free $k$ -pluriharmonic functions

- Let  $\mathbf{PH}^\infty(\mathbf{B}_n)$  be the vector space of all **bounded free  $k$ -pluriharmonic functions** on  $\mathbf{B}_n$ .

- For each  $m = 1, 2, \dots$ , define the norm  $\|\cdot\|_m : M_m(\mathbf{PH}^\infty(\mathbf{B}_n)) \rightarrow [0, \infty)$  by setting

$$\|[F_{ij}]_m\|_m := \sup \| [F_{ij}(\mathbf{X})]_m \|,$$

where sup is taken over all  $\mathbf{X} \in \mathbf{B}_n(\mathcal{H})$  and any  $\mathcal{H}$ .

- The norms  $\|\cdot\|_m$  determine an operator space structure on  $\mathbf{PH}^\infty(\mathbf{B}_n)$ , in the sense of **Ruan**.

# Bounded free $k$ -pluriharmonic functions

## Theorem

If  $F : \mathbf{B}_n(\mathcal{H}) \rightarrow B(\mathcal{H})$  is a free  $k$ -pluriharmonic function, then the  $F$  is bounded if and only if there exists  $A \in \mathcal{T}_n$  such that

$$F(\mathbf{X}) = \mathcal{B}_{\mathbf{X}}[A] := \mathbf{K}_{\mathbf{X}}^*(A \otimes I_{\mathcal{H}})\mathbf{K}_{\mathbf{X}}, \quad \mathbf{X} \in \mathbf{B}_n(\mathcal{H}).$$

In this case,  $A = \text{SOT-} \lim_{r \rightarrow 1} F(r\mathbf{S})$ .

Moreover, the map

$$\Phi : \mathbf{PH}^{\infty}(\mathbf{B}_n) \rightarrow \mathcal{T}_n \quad \text{defined by} \quad \Phi(F) := A$$

is a completely isometric isomorphism of operator spaces.

# Dirichlet extension problem for regular polyballs

- Let  $\mathbf{PH}^c(\mathbf{B}_n)$  be the set of all free  $k$ -pluriharmonic functions on  $\mathbf{B}_n$  which have continuous extensions to  $\mathbf{B}_n(\mathcal{H})^-$  (in norm topology), for any Hilbert space  $\mathcal{H}$ .
- Assume that  $\mathcal{H}$  is an infinite dimensional Hilbert space.



# Dirichlet extension problem for regular polyballs

## Theorem

If  $F : \mathbf{B}_n(\mathcal{H}) \rightarrow B(\mathcal{H})$  is a free  $k$ -pluriharmonic function, then  $F$  has a continuous extension to the closed polyball  $\mathbf{B}_n(\mathcal{H})^-$  (in the operator norm) if and only if there exists  $A \in \mathcal{P} := \text{span}\{f^*g : f, g \in \mathcal{A}_n\}^{-\|\cdot\|}$  such that

$$F(\mathbf{X}) = \mathcal{B}_{\mathbf{X}}[A], \quad \mathbf{X} \in \mathbf{B}_n(\mathcal{H}).$$

In this case,  $A = \lim_{r \rightarrow 1} F(r\mathbf{S})$ , where the convergence is in the operator norm. Moreover, the map

$$\Phi : \mathbf{PH}^c(\mathbf{B}_n) \rightarrow \mathcal{P} \quad \text{defined by} \quad \Phi(F) := A$$

is a completely isometric isomorphism of operator spaces.

# Noncommutative Poisson transforms of c.b. maps

- Consider the operator system

$$\mathcal{R}_n^* \mathcal{R}_n := \text{span}\{\mathbf{R}_\alpha^* \mathbf{R}_\beta : \alpha, \beta \in \mathbb{F}_{n_1}^+ \times \cdots \times \mathbb{F}_{n_k}^+\},$$

where  $\mathbf{R} := (\mathbf{R}_1, \dots, \mathbf{R}_k)$  and  $\mathbf{R}_i := (\mathbf{R}_{i,1}, \dots, \mathbf{R}_{i,n_i})$ .

- If  $\mu : \mathcal{R}_n^* \mathcal{R}_n \rightarrow B(\mathcal{E})$  is a completely bounded linear map, then there exists a unique completely bounded linear map

$$\widehat{\mu} := \mu \otimes id : \overline{\mathcal{R}_n^* \mathcal{R}_n}^{\|\cdot\|} \otimes_{\min} B(\mathcal{H}) \rightarrow B(\mathcal{E}) \otimes_{\min} B(\mathcal{H})$$

such that

$$\widehat{\mu}(A \otimes Y) = \mu(A) \otimes Y, \quad A \in \mathcal{R}_n^* \mathcal{R}_n, Y \in B(\mathcal{H}).$$

Moreover,  $\|\widehat{\mu}\|_{cb} = \|\mu\|_{cb}$  and, if  $\mu$  is completely positive, then so is  $\widehat{\mu}$ .

# Noncommutative Poisson transforms of c.b. maps

- Define the *free pluriharmonic Poisson kernel* by setting

$$\mathcal{P}(\mathbf{R}, \mathbf{X}) := \sum_{(\alpha, \beta) \in \Lambda} \mathbf{R}_{\tilde{\alpha}}^* \mathbf{R}_{\tilde{\beta}} \otimes \mathbf{X}_{\alpha} \mathbf{X}_{\beta}^*, \quad \mathbf{X} \in \mathbf{B}_n(\mathcal{H}),$$

where the convergence is in the operator norm topology, and  $(\alpha, \beta) \in \Lambda$  iff  $\alpha = (\alpha_1, \dots, \alpha_k)$  and  $\beta = (\beta_1, \dots, \beta_k)$ , with  $\alpha_i, \beta_i \in \mathbb{F}_{n_i}^+$ ,  $|\alpha_i| = m_i^-$ ,  $|\beta_i| = m_i^+$ .

- We introduce the *noncommutative Poisson transform of a c. b. map*  $\mu : \mathcal{R}_n^* \mathcal{R}_n \rightarrow B(\mathcal{E})$  on the regular polyball to be the map  $\mathcal{P}\mu : \mathbf{B}_n(\mathcal{H}) \rightarrow B(\mathcal{E}) \otimes_{\min} B(\mathcal{H})$  defined by

$$(\mathcal{P}\mu)(\mathbf{X}) := \widehat{\mu}[\mathcal{P}(\mathbf{R}, \mathbf{X})], \quad \mathbf{X} \in \mathbf{B}_n(\mathcal{H}).$$

# Noncommutative Poisson transforms of c.b. maps

## Theorem

Let  $\mu : \mathcal{R}_n^* \mathcal{R}_n \rightarrow B(\mathcal{E})$  be a completely bounded linear map. The following statements hold.

- (i) The map  $\mathbf{X} \mapsto \mathcal{P}(\mathbf{R}, \mathbf{X})$  is a positive  $k$ -pluriharmonic function on the polyball  $\mathbf{B}_n$ , with coefficients in  $B(\otimes_{i=1}^k F^2(H_{n_i}))$ , and has the factorization  $\mathcal{P}(\mathbf{R}, \mathbf{X}) = C_{\mathbf{X}}^* C_{\mathbf{X}}$ , where

$$C_{\mathbf{X}} := (I \otimes \Delta_{\mathbf{X}}(I)^{1/2}) \prod_{i=1}^k (I - \mathbf{R}_{i,1} \otimes X_{i,1}^* - \cdots - \mathbf{R}_{i,n_i} \otimes X_{i,n_i}^*)^{-1}.$$

- (ii) The noncommutative Poisson transform  $\mathcal{P}\mu$  is a free  $k$ -pluriharmonic function on the regular polyball  $\mathbf{B}_n$ .

# Noncommutative Poisson transforms of c.b. maps

(iii) If  $\mu$  is a completely positive linear map, then  $\mathcal{P}\mu$  is a positive free  $k$ -pluriharmonic function on  $\mathbf{B}_n$ .

- Let  $F$  be a free  $k$ -pluriharmonic function on the polyball  $\mathbf{B}_n$ , with operator-valued coefficients in  $B(\mathcal{E})$ , and with representation

$$F(\mathbf{X}) = \sum_{m_1 \in \mathbb{Z}} \cdots \sum_{m_k \in \mathbb{Z}} \sum_{(\alpha, \beta) \in \Lambda} A_{\alpha, \beta} \otimes \mathbf{X}_\alpha \mathbf{X}_\beta^*.$$

- We associate with  $F$  and each  $r \in [0, 1)$  the linear map  $\nu_{F_r} : \mathcal{R}_n^* \mathcal{R}_n \rightarrow B(\mathcal{E})$  by setting

$$\nu_{F_r}(\mathbf{R}_\alpha^* \tilde{\mathbf{R}}_\beta) := r^{|\alpha|+|\beta|} A_{\alpha, \beta}, \quad (\alpha, \beta) \in \Lambda.$$

# Noncommutative Poisson transforms of c.b. maps

## Theorem

Let  $F : \mathbf{B}_n(\mathcal{H}) \rightarrow B(\mathcal{E}) \otimes_{\min} B(\mathcal{H})$  be a free  $k$ -pluriharmonic function. Then the following statements are equivalent :

- (i) there exists a completely bounded linear map  $\mu : C^*(\mathbf{R}) \rightarrow B(\mathcal{E})$  such that  $F = \mathcal{P}\mu$ ;
- (ii) the linear maps  $\{\nu_{F_r}\}_{r \in [0,1]}$  associate with  $F$  are completely bounded and  $\sup_{0 \leq r < 1} \|\nu_{F_r}\|_{cb} < \infty$ ;

# Noncommutative Poisson transforms of c.b. maps

- (iii) there exists a  $k$ -tuple  $\mathbf{V} = (V_1, \dots, V_k)$  of doubly commuting row isometries acting on  $\mathcal{K}$  and bounded linear operators  $W_1, W_2 : \mathcal{E} \rightarrow \mathcal{K}$  such that

$$F(\mathbf{X}) = (W_1^* \otimes I) [C_{\mathbf{X}}(\mathbf{V})^* C_{\mathbf{X}}(\mathbf{V})] (W_2 \otimes I),$$

where

$$C_{\mathbf{X}}(\mathbf{V}) := (I \otimes \Delta_{\mathbf{X}}(I)^{1/2}) \prod_{i=1}^k (I - V_{i,1} \otimes X_{i,1}^* - \dots - V_{i,n_i} \otimes X_{i,n_i}^*)^{-1}.$$

Moreover, in this case we can choose  $\mu$  such that

$$\|\mu\|_{cb} = \sup_{0 \leq r < 1} \|\nu_{F_r}\|_{cb}.$$

# Noncommutative Poisson transforms of c.p. maps

## Corollary

Let  $F : \mathbf{B}_n(\mathcal{H}) \rightarrow B(\mathcal{E}) \otimes_{\min} B(\mathcal{H})$  be a free  $k$ -pluriharmonic function. Then the following statements are equivalent :

- (i) there exists a completely positive linear map  $\mu : C^*(\mathbf{R}) \rightarrow B(\mathcal{E})$  such that  $F = \mathcal{P}\mu$ ;
- (ii) the linear maps  $\{\nu_{F_r}\}_{r \in [0,1]}$  associate with  $F$  are completely positive ;
- (iii) there exists a  $k$ -tuple  $\mathbf{V} = (V_1, \dots, V_k)$  of doubly commuting row isometries acting on a Hilbert space  $\mathcal{K} \supset \mathcal{E}$  and a bounded operator  $W : \mathcal{E} \rightarrow \mathcal{K}$  such that

$$F(\mathbf{X}) = (W^* \otimes I) [C_{\mathbf{X}}(\mathbf{V})^* C_{\mathbf{X}}(\mathbf{V})] (W \otimes I).$$



# Noncommutative Poisson transforms of c.p. maps

- **Classical result** : A map  $u : \mathbb{D}^k \rightarrow \mathbb{C}$  is a positive  $k$ -harmonic function if and only if there is a finite positive Borel measure on  $\mathbb{T}^k$  such that

$$u(z) = \int_{\mathbb{T}^k} P(z, \zeta) d\mu(\zeta), \quad z \in \mathbb{D}^k,$$

where  $P(z, \zeta)$  is the Poisson kernel for the polydisk.

- **Open question** : Is any positive free  $k$ -pluriharmonic function on the regular polyball  $\mathbf{B}_n$  the noncommutative Poisson transform of a completely positive linear map  $\mu : C^*(\mathbf{R}) \rightarrow B(\mathcal{E})$ ?

# Noncommutative Poisson transforms of c.p. maps

- The answer is positive for the unit ball  $[B(\mathcal{H})^n]_1$  (when  $k = 1$ ) (P., Adv. Math., 2009) and for the regular polydisk  $\mathbf{D}^k(\mathcal{H})$  (when  $n_1 = \dots = n_k = 1$ ).

## Theorem

A map  $f : \mathbf{D}^k(\mathcal{H}) \rightarrow B(\mathcal{E}) \otimes_{\min} B(\mathcal{H})$  is a positive free  $k$ -pluriharmonic function on the **regular polydisk** if and only if there exists a completely positive linear map  $\mu : C^*(M_{Z_1}, \dots, M_{Z_k}) \rightarrow B(\mathcal{E})$  such that  $F = \mathcal{P}\mu$ , where  $M_{Z_1}, \dots, M_{Z_k}$  are the multiplication operators on  $H^2(\mathbb{D}^k)$ .

## Poincaré distance on the open unit disc

- The **hyperbolic (Poincaré) distance** on the open unit disc  $\mathbb{D} := \{z \in \mathbb{C} : |z| < 1\}$  is defined by

$$\delta_P(z, w) := \frac{1}{2} \ln \frac{1 + |\varphi_z(w)|}{1 - |\varphi_z(w)|}, \quad z, w \in \mathbb{D},$$

where  $\varphi_z$  is the automorphism of  $\mathbb{D}$  given by  $\varphi_z(w) = \frac{w-z}{1-\bar{z}w}$ .

# Poincaré distance on the open unit disc

- **Basic properties of the Poincaré distance :**

- 1 the Poincaré distance is invariant under the conformal automorphisms of  $\mathbb{D}$ , i.e.,

$$\delta_P(\varphi(z), \varphi(w)) = \delta_P(z, w), \quad z, w \in \mathbb{D},$$

for all  $\varphi \in \text{Aut}(\mathbb{D})$ ;

- 2 the  $\delta_P$ -topology induced on the open disc is the usual planar topology;
- 3  $(\mathbb{D}, \delta_P)$  is a complete metric space;
- 4 any analytic function  $f : \mathbb{D} \rightarrow \mathbb{D}$  is distance-decreasing, i.e.,

$$\delta_P(f(z), f(w)) \leq \delta_P(z, w), \quad z, w \in \mathbb{D}.$$

## Extensions of Poincaré distance

- **Bergman** introduced an analogue of the Poincaré distance for the open unit ball of  $\mathbb{C}^n$ ,

$$\mathbb{B}_n := \{z = (z_1, \dots, z_n) \in \mathbb{C}^n : \|z\|_2 < 1\},$$

defined by

$$\beta_n(z, w) = \frac{1}{2} \ln \frac{1 + \|\psi_z(w)\|_2}{1 - \|\psi_z(w)\|_2}, \quad z, w \in \mathbb{B}_n,$$

where  $\psi_z$  is the involutive automorphism of  $\mathbb{B}_n$  that interchanges 0 and  $z$ . The **Poincaré-Bergman distance** has properties similar to those of  $\delta_P$ .

# Extensions of Poincaré distance

- There are several extensions of the Poincaré-Bergman distance to more general domains.
  - 1 The work of R.S. Phillips and L. Harris on infinite-dimensional Cartan domains.
  - 2 The work of Suciu, Foiaş, and Andô-Suciu-Timotin on Harnack type distances between two contractions.
  - 3 The work of P. on hyperbolic geometry on  $[B(\mathcal{H})^n]_1$ .

# Harnack domination

- Preorder relation  $\overset{H}{\prec}$  on the closed ball  $\mathbf{B}_n(\mathcal{H})^-$ .

## Definition

If  $\mathbf{A}$  and  $\mathbf{B}$  are in  $\mathbf{B}_n(\mathcal{H})^-$ , we say that  $\mathbf{A}$  is *Harnack dominated* by  $\mathbf{B}$ , and denote  $\mathbf{A} \overset{H}{\prec} \mathbf{B}$ , if there exists  $c > 0$  such that

$$F(r\mathbf{A}) \leq c^2 F(r\mathbf{B})$$

for any positive free  $k$ -pluriharmonic function  $F$  with operator valued coefficients and any  $r \in [0, 1)$ . When we want to emphasize the constant  $c$ , we write  $\mathbf{A} \overset{H}{\prec}_c \mathbf{B}$ .

# Harnack equivalence

## Definition

If  $\mathbf{A}, \mathbf{B} \in \mathbf{B}_n(\mathcal{H})^-$ , we say that  $\mathbf{A}$  and  $\mathbf{B}$  are *Harnack equivalent* (and denote  $\mathbf{A} \stackrel{H}{\sim} \mathbf{B}$ ) if there exists  $c > 1$  such that

$$\frac{1}{c^2}F(r\mathbf{B}) \leq F(r\mathbf{A}) \leq c^2F(r\mathbf{B}), \quad r \in [0, 1),$$

for any positive free  $k$ -pluriharmonic function

$F : \mathbf{B}_n(\mathcal{H}) \rightarrow B(\mathcal{E}) \otimes_{\min} B(\mathcal{H})$ , where  $\mathcal{E}$  is a separable Hilbert space. In this case, we write  $\mathbf{A} \stackrel{H}{\sim}_c \mathbf{B}$ .

- The equivalence classes with respect to the equivalence relation  $\stackrel{H}{\sim}$  are called **Harnack parts** of  $\mathbf{B}_n(\mathcal{H})^-$ .



## Poisson domination

- Recall the *free pluriharmonic Poisson kernel* :

$$\mathcal{P}(\mathbf{R}, \mathbf{X}) := \sum_{(\alpha, \beta) \in \Lambda} \mathbf{R}_{\tilde{\alpha}}^* \mathbf{R}_{\tilde{\beta}} \otimes \mathbf{X}_{\alpha} \mathbf{X}_{\beta}^*$$

for any  $\mathbf{X} \in \mathbf{B}_n(\mathcal{H})$ , where the convergence is in the operator norm topology.

- If  $\mathbf{A}$  and  $\mathbf{B}$  are in  $\mathbf{B}_n(\mathcal{H})^-$ , we say that  $\mathbf{A}$  is *Poisson dominated* by  $\mathbf{B}$ , and denote  $\mathbf{A} \stackrel{P}{\prec} \mathbf{B}$ , if there exists  $c > 0$  such that

$$\mathcal{P}(\mathbf{R}, r\mathbf{A}) \leq c^2 \mathcal{P}(\mathbf{R}, r\mathbf{B})$$

for any  $r \in [0, 1)$ . When we want to emphasize the constant  $c$ , we write  $\mathbf{A} \stackrel{P}{\prec}_c \mathbf{B}$ .

# Poisson equivalence

## Definition

If  $\mathbf{A}, \mathbf{B} \in \mathbf{B}_n(\mathcal{H})^-$ , we say that  $\mathbf{A}$  and  $\mathbf{B}$  are **Poisson equivalent** (we denote  $\mathbf{A} \stackrel{P}{\sim} \mathbf{B}$ ) if and only if there exists  $c \geq 1$  such that

$$\frac{1}{c^2} \mathcal{P}(\mathbf{R}, r\mathbf{B}) \leq \mathcal{P}(\mathbf{R}, r\mathbf{A}) \leq c^2 \mathcal{P}(\mathbf{R}, r\mathbf{B})$$

for any  $r \in [0, 1)$ .

We also use the notation  $\mathbf{A} \underset{c}{\stackrel{P}{\sim}} \mathbf{B}$  if  $\mathbf{A} \underset{c}{\prec} \mathbf{B}$  and  $\mathbf{B} \underset{c}{\prec} \mathbf{A}$ .

# Harnack inequality

## Theorem

Let  $F$  be a positive free  $k$ -pluriharmonic function on the regular polyball  $\mathbf{B}_n$ , with operator coefficients in  $B(\mathcal{E})$  and let  $0 \leq r < 1$ . Then

$$F(0) \left( \frac{1-r}{1+r} \right)^k \leq F(\mathbf{X}) \leq F(0) \left( \frac{1+r}{1-r} \right)^k$$

for any  $\mathbf{X} \in r\mathbf{B}_n(\mathcal{H})^-$ .

# Harnack and Poisson equivalence class containing 0

## Theorem

Let  $\mathbf{A} = (A_1, \dots, A_k) \in \mathbf{B}_n(\mathcal{H})^-$ . Then the following statements are equivalent.

- 1  $\mathbf{A} \stackrel{H}{\sim} 0$ ;
- 2  $r(A_i) < 1$  for any  $i \in \{1, \dots, k\}$  and there exists  $a > 0$  such that

$$\mathcal{P}(\mathbf{R}, r\mathbf{A}) \geq aI, \quad r \in [0, 1);$$

- 3  $\mathbf{A} \in \mathbf{B}_n(\mathcal{H})$ ;
- 4  $\mathbf{A} \stackrel{P}{\sim} 0$ .

## Hyperbolic metric on Harnack parts

- Given  $\mathbf{A}, \mathbf{B} \in \mathbf{B}_n(\mathcal{H})^-$  in the same Harnack part, i.e.  $\mathbf{A} \stackrel{H}{\sim} \mathbf{B}$ , we introduce

$$\omega_H(\mathbf{A}, \mathbf{B}) := \inf \left\{ c > 1 : \mathbf{A} \stackrel{H}{\sim}_c \mathbf{B} \right\}.$$

### Theorem

Let  $\Delta$  be a Harnack part of  $\mathbf{B}_n(\mathcal{H})^-$  and define  $\delta_H : \Delta \times \Delta \rightarrow \mathbb{R}^+$  by setting

$$\delta_H(\mathbf{A}, \mathbf{B}) := \ln \omega_H(\mathbf{A}, \mathbf{B}), \quad \mathbf{A}, \mathbf{B} \in \Delta.$$

Then  $\delta_H$  is a metric on  $\Delta$ .

## Hyperbolic metric on Harnack parts

- **Schwarz-Pick lemma** for free holomorphic functions on the regular polyball  $\mathbf{B}_n$  with operator-valued coefficients, with respect to the hyperbolic metric.

### Theorem

Let  $\Phi = (\Phi_1, \dots, \Phi_m) : \mathbf{B}_n(\mathcal{H}) \rightarrow [B(\mathcal{H})^m]_1^-$  be a free holomorphic function on the regular polyball. If  $\mathbf{X}, \mathbf{Y} \in \mathbf{B}_n(\mathcal{H})$ , then  $\Phi(\mathbf{X}) \stackrel{H}{\sim} \Phi(\mathbf{Y})$  and

$$\delta_H(\Phi(\mathbf{X}), \Phi(\mathbf{Y})) \leq \delta_H(\mathbf{X}, \mathbf{Y}),$$

where  $\delta_H$  is the hyperbolic metric defined on the Harnack parts of  $[B(\mathcal{H})^m]_1^-$  and on the polyball  $\mathbf{B}_n(\mathcal{H})$ , respectively.

# Hyperbolic metric on Harnack parts

- The hyperbolic metric is invariant under the group  $Aut(\mathbf{B}_n)$  of all free holomorphic automorphisms of  $\mathbf{B}_n$ .

## Theorem

Let  $\mathbf{A}$  and  $\mathbf{B}$  be in  $\mathbf{B}_n(\mathcal{H})^-$  such that  $\mathbf{A} \stackrel{H}{\sim} \mathbf{B}$ . Then

$$\delta_H(\mathbf{A}, \mathbf{B}) = \delta_H(\Psi(\mathbf{A}), \Psi(\mathbf{B})), \quad \Psi \in Aut(\mathbf{B}_n).$$

## Metric on Poisson parts of the polyball

- Given  $\mathbf{A}, \mathbf{B} \in \mathbf{B}_n(\mathcal{H})^-$  in the same Poisson part, i.e.  $\mathbf{A} \overset{P}{\sim} \mathbf{B}$ , we introduce

$$\omega_{\mathcal{P}}(\mathbf{A}, \mathbf{B}) := \inf \left\{ c > 1 : \mathbf{A} \overset{P}{\sim}_c \mathbf{B} \right\}.$$

### Theorem

Let  $\Delta$  be a Poisson part of  $\mathbf{B}_n(\mathcal{H})^-$  and define the function  $\delta_{\mathcal{P}} : \Delta \times \Delta \rightarrow \mathbb{R}^+$  by setting

$$\delta_{\mathcal{P}}(\mathbf{A}, \mathbf{B}) := \ln \omega_{\mathcal{P}}(\mathbf{A}, \mathbf{B}), \quad \mathbf{A}, \mathbf{B} \in \Delta.$$

Then  $\delta_{\mathcal{P}}$  is a metric on  $\Delta$ .



# Metric on Poisson parts of the polyball

## Theorem

If  $\mathbf{A}$  and  $\mathbf{B}$  are in the open ball  $\mathbf{B}_n(\mathcal{H})$ , then

$$\delta_{\mathcal{P}}(\mathbf{A}, \mathbf{B}) = \ln \max \left\{ \left\| C_{\mathbf{A}}(\mathbf{R}) C_{\mathbf{B}}(\mathbf{R})^{-1} \right\|, \left\| C_{\mathbf{B}}(\mathbf{R}) C_{\mathbf{A}}(\mathbf{R})^{-1} \right\| \right\},$$

where

$$C_{\mathbf{X}}(\mathbf{R}) := (I \otimes \Delta_{\mathbf{X}}(I)^{1/2}) \prod_{i=1}^k (I - \mathbf{R}_{i,1} \otimes X_{i,1}^* - \cdots - \mathbf{R}_{i,n_i} \otimes X_{i,n_i}^*)^{-1}$$

for any  $\mathbf{X} = (X_1, \dots, X_k) \in \mathbf{B}_n(\mathcal{H})$  with  $X_i = (X_{i,1}, \dots, X_{i,n_i})$ .

## Metric on Poisson parts of the polyball

- Set

$$\mathbf{B}_n(\mathcal{H})_0^- := \left\{ \mathbf{X} \in \mathbf{B}_n(\mathcal{H})^- : \mathbf{X} \stackrel{P}{\prec} 0 \right\}$$

and recall that  $\mathbf{B}_n(\mathcal{H}) \subset \mathbf{B}_n(\mathcal{H})_0^-$ .

### Theorem

*Let  $\Delta$  be a Poisson part of  $\mathbf{B}_n(\mathcal{H})_0^-$ . Then the following properties hold :*

- (i)  $\delta_{\mathcal{P}}$  is a complete metric on  $\Delta$ .
- (ii) the  $\delta_{\mathcal{P}}$ -topology and the operator norm topology coincide on the open polyball  $\mathbf{B}_n(\mathcal{H})$ .
- (iii) the  $\delta_H$ -topology is stronger than the  $\delta_{\mathcal{P}}$ -topology on  $\mathbf{B}_n(\mathcal{H})$ .

# Positive $k$ -harmonic functions on the regular polydisk

## Theorem

Let  $F : \mathbf{D}^k(\mathcal{H}) \rightarrow B(\mathcal{E}) \otimes_{\min} B(\mathcal{H})$  be a free  $k$ -pluriharmonic function. Then the following statements are equivalent :

- (i)  $F$  is positive ;
- (ii) there exists a completely positive linear map  $\mu : C^*(\mathbf{R}) \rightarrow B(\mathcal{E})$  such that  $F = \mathcal{P}\mu$  ;
- (iii) there exists a  $k$ -tuple  $\mathbf{U} = (U_1, \dots, U_k)$  of commuting unitaries acting on a Hilbert space  $\mathcal{K} \supset \mathcal{E}$  and a bounded operator  $W : \mathcal{E} \rightarrow \mathcal{K}$  such that

$$F(\mathbf{X}) = (W^* \otimes I) [C_{\mathbf{X}}(\mathbf{U})^* C_{\mathbf{X}}(\mathbf{U})] (W \otimes I),$$

# Positive $k$ -harmonic functions on the regular polydisk

where

$$C_{\mathbf{X}}(\mathbf{U}) := (I \otimes \Delta_{\mathbf{X}}(I)^{1/2}) \prod_{i=1}^k (I - U_i \otimes X_i^*)$$

for any  $\mathbf{X} = (X_1, \dots, X_k) \in \mathbf{D}^k(\mathcal{H})$ .

- The **Kobayashi distance** for the polydisc  $\mathbb{D}^k$  is given by

$$K_{\mathbb{D}^k}(\mathbf{z}, \mathbf{w}) = \frac{1}{2} \ln \frac{1 + \|\psi_{\mathbf{z}}(\mathbf{w})\|_{\infty}}{1 - \|\psi_{\mathbf{z}}(\mathbf{w})\|_{\infty}},$$

where  $\psi_{\mathbf{z}}$  is the involutive automorphisms of  $\mathbb{D}^k$  given by

$$\psi_{\mathbf{z}} = \left( \frac{w_1 - z_1}{1 - \bar{z}_1 w_1}, \dots, \frac{w_k - z_k}{1 - \bar{z}_k w_k} \right)$$

for any  $\mathbf{z} = (z_1, \dots, z_k)$  and  $\mathbf{w} = (w_1, \dots, w_k)$  in  $\mathbb{D}^k$ .

# Hyperbolic metric on the regular polydisk

## Theorem

Let  $\mathbf{D}^k(\mathcal{H})$  be the regular polydisk. The following statements hold.

- (i) If  $\mathbf{A}, \mathbf{B} \in \mathbf{D}^k(\mathcal{H})^-$ , then  $\mathbf{A} \stackrel{H}{\sim} \mathbf{B}$  if and only if  $\mathbf{A} \stackrel{P}{\sim} \mathbf{B}$ .
- (ii) The metrics  $\delta_H$  and  $\delta_P$  coincide on the Harnack parts of  $\mathbf{D}^k(\mathcal{H})^-$ .
- (iii) If  $\mathbf{A}$  and  $\mathbf{B}$  are in  $\mathbf{D}^k(\mathcal{H})^-$  and  $\mathbf{A} \stackrel{H}{\sim} \mathbf{B}$ , then

$$\delta_H(\mathbf{A}, \mathbf{B}) = \delta_H(\Psi(\mathbf{A}), \Psi(\mathbf{B})), \quad \Psi \in \text{Aut}(\mathbf{D}^k).$$

## Hyperbolic metric on the regular polydisk

(iv) If  $\mathbf{A}$  and  $\mathbf{B}$  are in  $\mathbf{D}^k(\mathcal{H})$ , then

$$\delta_H(\mathbf{A}, \mathbf{B}) = \ln \max \left\{ \left\| C_{\mathbf{A}}(\mathbf{R}) C_{\mathbf{B}}(\mathbf{R})^{-1} \right\|, \left\| C_{\mathbf{B}}(\mathbf{R}) C_{\mathbf{A}}(\mathbf{R})^{-1} \right\| \right\},$$

where

$$C_{\mathbf{X}}(\mathbf{R}) := (I \otimes \Delta_{\mathbf{X}}(I)^{1/2}) \prod_{i=1}^k (I - R_i \otimes X_i^*)$$

for any  $\mathbf{X} = (X_1, \dots, X_k) \in \mathbf{D}^k(\mathcal{H})$ .

## Hyperbolic metric on the regular polydisk

- (v)  $\delta_H|_{\mathbb{D}^k \times \mathbb{D}^k}$  is equivalent to the Kobayashi distance on the polydisk  $\mathbb{D}^k$  and

$$\delta_H(\mathbf{z}, \mathbf{w}) = \frac{1}{2} \ln \frac{\prod_{i=1}^k (1 + |\psi_{z_i}(\mathbf{w}_i)|)}{\prod_{i=1}^k (1 - |\psi_{z_i}(\mathbf{w}_i)|)}$$

for any  $\mathbf{z} = (z_1, \dots, z_k)$  and  $\mathbf{w} = (w_1, \dots, w_k)$  in  $\mathbb{D}^k$ , where  $\psi_{\mathbf{z}} := (\psi_{z_1}, \dots, \psi_{z_n})$  is the involutive automorphisms of  $\mathbb{D}^k$  such that  $\psi_{z_i}(0) = z_i$  and  $\psi_{z_i}(z_i) = 0$ .

- (vi) The hyperbolic metric  $\delta_H$  is complete on the Harnack parts of  $\mathbf{D}^k(\mathcal{H})_0^-$ .
- (vii) The  $\delta_H$ -topology coincides with the operator norm topology on the regular polydisk  $\mathbf{D}^k(\mathcal{H})$ .

# Hyperbolic metric on the regular polydisk

## Corollary

Let  $f = (f_1, \dots, f_m) : \mathbf{D}^k(\mathcal{H}) \rightarrow [B(\mathcal{H})^m]_1$  be a free holomorphic function on the regular polydisk. If  $\mathbf{X}, \mathbf{Y} \in \mathbf{D}^k(\mathcal{H})$ , then

$$\delta_H(f(\mathbf{X}), f(\mathbf{Y})) \leq \delta_H(\mathbf{X}, \mathbf{Y}),$$

where  $\delta_H$  is the hyperbolic metric. In particular, if  $f(0) = 0$ , then

$$\frac{1 + \|f(\mathbf{z})\|_2}{1 - \|f(\mathbf{z})\|_2} \leq \prod_{i=1}^k \frac{1 + |z_i|}{1 - |z_i|}$$

for any  $\mathbf{z} = (z_1, \dots, z_k)$  in  $\mathbb{D}^k$ .



# Herglotz-Riesz representations

- Define the space

$$\mathbf{RH}(\mathbf{B}_n) := \text{span} \{ \mathfrak{R}f : f \in \text{Hol}_{\mathcal{E}}(\mathbf{B}_n) \},$$

where  $\text{Hol}_{\mathcal{E}}(\mathbf{B}_n)$  is the set of all free holomorphic functions in the polyball  $\mathbf{B}_n$ , with coefficients in  $B(\mathcal{E})$ .

- If  $\varphi \in \mathbf{RH}(\mathbf{B}_n)$ , we consider the family  $\{\nu_{\varphi_r}\}_{r \in [0,1]}$  of linear maps  $\nu_{\varphi_r} : \mathcal{R}_n^* \mathcal{R}_n \rightarrow B(\mathcal{E})$ . Note that  $\nu_{\varphi_r}(\mathbf{R}_{\alpha}^* \mathbf{R}_{\beta}) = 0$  if  $\mathbf{R}_{\alpha}^* \mathbf{R}_{\beta}$  is different from  $\mathbf{R}_{\gamma}$  or  $\mathbf{R}_{\gamma}^*$  for some  $\gamma \in \mathbf{F}_n^+$ .

# Herglotz-Riesz representations

- Let  $\mu : \mathcal{R}_n^* \mathcal{R}_n \rightarrow B(\mathcal{E})$  be a completely positive linear map. The *noncommutative Herglotz-Riesz transform* of  $\mu$  on the regular polyball is the map  $\mathbf{H}\mu : \mathbf{B}_n(\mathcal{H}) \rightarrow B(\mathcal{E}) \otimes_{\min} B(\mathcal{H})$  defined by

$$(\mathbf{H}\mu)(\mathbf{X}) := \widehat{\mu} \left[ 2 \prod_{i=1}^k (I - \mathbf{R}_{i,1}^* \otimes X_{i,1} - \dots - \mathbf{R}_{i,n_i}^* \otimes X_{i,n_i})^{-1} - I \right]$$

for  $\mathbf{X} := (X_1, \dots, X_k) \in \mathbf{B}_n(\mathcal{H})$ .

# Herglotz-Riesz representations

## Theorem

Let  $f$  be a free holomorphic function from the polyball  $\mathbf{B}_n(\mathcal{H})$  to  $B(\mathcal{E}) \otimes_{\min} B(\mathcal{H})$ . Then the following statements are equivalent.

- (i)  $f$  is a free holomorphic function with  $\Re f \geq 0$  and the linear maps  $\{\nu_{\Re f_r}\}_{r \in [0,1]}$  associated with  $\Re f$  are completely positive.
- (ii) The function  $f$  admits a **Herglotz-Riesz representation**

$$f(\mathbf{X}) = (\mathbf{H}\mu)(\mathbf{X}) + i\Im f(0),$$

where  $\mu : C^*(\mathbf{R}) \rightarrow B(\mathcal{E})$  is a completely positive linear map with the property that  $\mu(\mathbf{R}_\alpha^* \mathbf{R}_\beta) = 0$  if  $\mathbf{R}_\alpha^* \mathbf{R}_\beta$  is not equal to  $\mathbf{R}_\gamma$  or  $\mathbf{R}_\gamma^*$  for some  $\gamma \in \mathbf{F}_n^+$ .

# Herglotz-Riesz representations

- (iii) There exist a  $k$ -tuple  $\mathbf{V} = (V_1, \dots, V_k)$  of doubly commuting row isometries on a Hilbert space  $\mathcal{K}$ , and a bounded linear operator  $W : \mathcal{E} \rightarrow \mathcal{K}$ , such that

$$f(\mathbf{X}) = (W^* \otimes I) \left[ 2 \prod_{i=1}^k (I - V_{i,1}^* \otimes X_{i,1} - \dots - V_{i,n_i}^* \otimes X_{i,n_i})^{-1} - I \right] \times (W \otimes I) + i\Im f(0)$$

and  $W^* \mathbf{V}_\alpha^* \mathbf{V}_\beta W = 0$  if  $\mathbf{R}_\alpha^* \mathbf{R}_\beta$  is not equal to  $\mathbf{R}_\gamma$  or  $\mathbf{R}_\gamma^*$  for some  $\gamma \in \mathbf{F}_n^+$ .

# Herglotz-Riesz representations

- When  $n_1 = \dots = n_k = 1$ , we obtain an operator-valued extension of **Korányi-Pukánszky** integral representation.

## Theorem

*If  $n_1 = \dots = n_k = 1$ , then the statements in the theorem above are equivalent to*

- (iv) *The map  $f : \mathbf{D}^k(\mathcal{H}) \rightarrow B(\mathcal{E}) \otimes_{\min} B(\mathcal{H})$  is a free holomorphic function on the regular polydisk and  $\Re f \geq 0$ .*

# Herglotz-Riesz representations

- **Korányi-Pukánzky result :**

## Theorem

*A function  $f : \mathbb{D}^k \rightarrow \mathbb{C}$  is holomorphic and  $\Re f \geq 0$  if and only if it admits a representation*

$$f(z) = i\Im f(0) + \int_{\mathbb{T}^k} \left[ 2 \prod_{j=1}^k \frac{1}{1 - z_j \bar{\zeta}_j} - 1 \right] d\mu(\zeta)$$

*where  $\mu$  is a positive measure on  $\mathbb{T}^k$  such that, unless  $m_j \geq 0$  for any  $j \in \{1, \dots, k\}$  or  $m_k \leq 0$  for any  $j \in \{1, \dots, k\}$ ,*

$$\int_{\mathbb{T}^k} \zeta_1^{m_1} \cdots \zeta_k^{m_k} d\mu(\zeta) = 0.$$

**THANK YOU**

## Naimark dilations

- We provide a **Naimark type dilation** theorem for direct products  $\mathbb{F}_{n_1}^+ \times \cdots \times \mathbb{F}_{n_k}^+$  of unital free semigroups and characterize the positive free  $k$ -pluriharmonic functions.
- Let  $\mathbf{F}_n^+ := \mathbb{F}_{n_1}^+ \times \cdots \times \mathbb{F}_{n_k}^+$  be the unital semigroup with neutral element  $\mathbf{g} := (g_0^1, \dots, g_0^k)$ .
- Let  $\omega = (\omega_1, \dots, \omega_k)$ ,  $\gamma = (\gamma_1, \dots, \gamma_k)$ ,  $\alpha := (\alpha_1, \dots, \alpha_k)$ , and  $\beta := (\beta_1, \dots, \beta_k)$  be in  $\mathbf{F}_n^+$ .

### Definition

We say that  $K : \mathbf{F}_n^+ \times \mathbf{F}_n^+ \rightarrow B(\mathcal{E})$  is a **left  $k$ -multi-Toeplitz kernel** if  $K(\mathbf{g}, \mathbf{g}) = I_{\mathcal{E}}$  and

$$K(\sigma, \omega) = \begin{cases} K(\alpha, \beta) & \text{if } \mathbf{S}_{\sigma}^* \mathbf{S}_{\omega} = \mathbf{S}_{\alpha}^* \mathbf{S}_{\beta} \\ 0 & \text{if } \mathbf{S}_{\sigma}^* \mathbf{S}_{\omega} = 0. \end{cases}$$



## Naimark dilations

- We say that  $\Gamma : \mathbf{F}_n^+ \times \mathbf{F}_n^+ \rightarrow B(\mathcal{E})$  is a **right  $k$ -multi-Toeplitz kernel** if  $\Gamma(\tilde{\sigma}, \tilde{\omega}) = K(\sigma, \omega)$ , where  $\tilde{\sigma} = (\tilde{\sigma}_1, \dots, \tilde{\sigma}_k)$  and  $\tilde{\sigma}_i := g_{j_m}^i \cdots g_{j_1}^i$  is the reverse of  $\sigma_i := g_{j_1}^i \cdots g_{j_m}^i$ .

### Definition

A map  $K : \mathbf{F}_n^+ \times \mathbf{F}_n^+ \rightarrow B(\mathcal{E})$  has a **Naimark dilation** if there exists a  $k$ -tuple of commuting row isometries  $\mathbf{V} = (V_1, \dots, V_k)$ ,  $V_i = (V_{i,1}, \dots, V_{i,n_i})$ , on a Hilbert space  $\mathcal{K} \supset \mathcal{E}$ , i.e. the non-selfadjoint algebra  $Alg(V_i)$  commutes with  $Alg(V_s)$  for any  $i, s \in \{1, \dots, k\}$  with  $i \neq s$ , such that

$$K(\sigma, \omega) = P_{\mathcal{E}} \mathbf{V}_{\sigma}^* \mathbf{V}_{\omega} |_{\mathcal{E}}, \quad \sigma, \omega \in \mathbf{F}_n^+.$$

The dilation is called **minimal** if  $\mathcal{K} = \bigvee_{\omega \in \mathbf{F}_n^+} \mathbf{V}_{\omega} \mathcal{E}$ .

## Naimark dilations

### Theorem

*A map  $K : \mathbf{F}_n^+ \times \mathbf{F}_n^+ \rightarrow B(\mathcal{H})$  is a positive semi-definite left  $k$ -multi-Toeplitz kernel on  $\mathbf{F}_n^+$  if and only if it admits a Naimark dilation. In this case, there is a minimal dilation which is uniquely determined up to an isomorphism.*

### Theorem

*A map  $\Gamma : \mathbf{F}_n^+ \times \mathbf{F}_n^+ \rightarrow B(\mathcal{H})$  is a positive semi-definite right  $k$ -multi-Toeplitz kernel on  $\mathbf{F}_n^+$  if and only if it admits a Naimark dilation. In this case, there is a minimal dilation which is uniquely determined up to an isomorphism.*

## Schur type results

- If  $F$  is a free  $k$ -pluriharmonic function on the polyball  $\mathbf{B}_n$  with operator-valued coefficients in  $B(\mathcal{E})$ , one can associate a right  $k$ -multi-Toeplitz kernel  $\Gamma_F$  on  $\mathbf{F}_n^+$  in terms of the coefficients of  $F$ .
- **Schur type result** for positive  $k$ -pluriharmonic functions in polyballs.

### Theorem

*Let  $F$  be a  $k$ -pluriharmonic function on the regular polyball  $\mathbf{B}_n$ , with coefficients in  $B(\mathcal{E})$ . Then  $F$  is positive on  $\mathbf{B}_n$  if and only if the kernel  $\Gamma_{F_r}$  is positive semi-definite for any  $r \in [0, 1)$ , where  $F_r$  stands for the mapping  $\mathbf{X} \mapsto F(r\mathbf{X})$ .*

# Schur type results

## Definition

A free holomorphic function on the polyball  $\mathbf{B}_n$  and with operator-valued coefficients in  $B(\mathcal{E})$  has the form

$$f(\mathbf{X}) = \sum_{m_1 \in \mathbb{N}} \cdots \sum_{m_k \in \mathbb{N}} \sum_{\substack{\alpha_j \in \mathbb{F}_{n_j}^+, i \in \{1, \dots, k\} \\ |\alpha_j| = m_j}} A_{(\alpha_1, \dots, \alpha_k)} \otimes X_{1, \alpha_1} \cdots X_{k, \alpha_k},$$

where  $\mathbf{X} = (X_1, \dots, X_k) \in \mathbf{B}_n(\mathcal{H})$  and the series converge in the operator norm topology.

# Schur type results

## Corollary

Let  $f : \mathbf{B}_n(\mathcal{H}) \rightarrow B(\mathcal{E}) \otimes_{\min} B(\mathcal{H})$  be a free holomorphic function. Then the following statements are equivalent.

- (i)  $\Re f \geq 0$  on the polyball  $\mathbf{B}_n$ ;
- (ii)  $\Re f(r\mathbf{S}) \geq 0$  for any  $r \in [0, 1)$ ;
- (iii) the right  $k$ -multi Toeplitz kernel  $\Gamma_{\Re f_r}$  is positive semidefinite for any  $r \in [0, 1)$ .

# Positive $k$ -pluriharmonic functions

## Theorem

A map  $F : \mathbf{B}_n(\mathcal{H}) \rightarrow B(\mathcal{E}) \otimes_{\min} B(\mathcal{H})$ , with  $F(0) = I$ , is a positive free  $k$ -pluriharmonic function on the regular polyball if and only if it has the form

$$F(\mathbf{X}) = \sum_{(\alpha, \beta) \in \Lambda} P_{\mathcal{E}} \mathbf{V}_{\tilde{\alpha}}^* \mathbf{V}_{\tilde{\beta}}|_{\mathcal{E}} \otimes \mathbf{X}_{\alpha} \mathbf{X}_{\beta}^*,$$

where  $\mathbf{V} = (V_1, \dots, V_k)$  is a  $k$ -tuple of commuting row isometries on a space  $\mathcal{K} \supset \mathcal{E}$  such that

$$\sum_{(\alpha, \beta) \in \Lambda} P_{\mathcal{E}} \mathbf{V}_{\tilde{\alpha}}^* \mathbf{V}_{\tilde{\beta}}|_{\mathcal{E}} \otimes r^{|\alpha|+|\beta|} \mathbf{S}_{\alpha} \mathbf{S}_{\beta}^* \geq 0, \quad r \in [0, 1),$$

and the series is convergent in the operator topology.

# Positive $k$ -pluriharmonic functions

## Definition

A  $k$ -tuple  $\mathbf{V} = (V_1, \dots, V_k)$  of commuting row isometries  $V_i = (V_{i,1}, \dots, V_{i,n_i})$  is called *pluriharmonic* if the free  $k$ -pluriharmonic Poisson kernel

$$\mathcal{P}(\mathbf{V}, r\mathbf{S}) := \sum_{(\alpha, \beta) \in \Lambda} \mathbf{V}_{\tilde{\alpha}}^* \mathbf{V}_{\tilde{\beta}} \otimes r^{|\alpha|+|\beta|} \mathbf{S}_{\alpha} \mathbf{S}_{\beta}^*$$

is a positive operator for any  $r \in [0, 1)$ .

- Example :  $\mathbf{V} := \mathbf{R} = (\mathbf{R}_1, \dots, \mathbf{R}_k)$ .

# Positive $k$ -pluriharmonic functions

## Proposition

Let  $\mathbf{V} = (V_1, \dots, V_k)$ ,  $V_i = (V_{i,1}, \dots, V_{i,n_i})$ , be a  $k$ -tuple of commuting row isometries. Then  $\mathbf{V}$  is pluriharmonic in each of the following particular cases :

- (i) if  $k = 1$  and  $n_1 \in \mathbb{N}$ ;
- (ii) if  $\mathbf{V}$  is doubly commuting, i.e. the  $C^*$ -algebra  $C^*(V_i)$  commutes with  $C^*(V_s)$  if  $i, s \in \{1, \dots, k\}$  with  $i \neq s$ ;
- (iii) if  $n_1 = \dots = n_k = 1$ .



# Positive $k$ -pluriharmonic functions

## Proposition

Let  $\mathbf{V} = (V_1, \dots, V_k)$  be a pluriharmonic tuple of commuting row isometries on a Hilbert space  $\mathcal{K}$  and let  $\mathcal{E} \subset \mathcal{K}$  be a subspace. Then the map

$$F(\mathbf{X}) := (P_{\mathcal{E}} \otimes I) \mathcal{P}(\mathbf{V}, \mathbf{X})|_{\mathcal{E} \otimes \mathcal{H}}, \quad \mathbf{X} \in \mathbf{B}_n(\mathcal{H})$$

is a positive free  $k$ -pluriharmonic function on the polyball  $\mathbf{B}_n$  with operator-valued coefficients in  $B(\mathcal{E})$ , and  $F(0) = I$ .

Moreover, in the particular cases when  $k = 1$  (P., [Adv. Math., 2009](#)), or when  $n_1 = \dots = n_k = 1$ , each positive free  $k$ -pluriharmonic function  $F$  with  $F(0) = I$  has the form above.

## Positive $k$ -pluriharmonic functions

- In particular, we obtain a structure theorem for the positive  $k$ -harmonic functions on the regular polydisk  $\mathbf{D}^k(\mathcal{H})$ , which extends the corresponding classical result on scalar polydisks.

### Corollary

*A map  $F : \mathbf{D}^k(\mathcal{H}) \rightarrow B(\mathcal{E}) \otimes_{\min} B(\mathcal{H})$  is positive free  $k$ -pluriharmonic function with  $F(0) = I$  if and only if there is a  $k$ -tuple of doubly commuting isometries  $\mathbf{V} = (V_1, \dots, V_k)$  on a Hilbert space  $\mathcal{K} \supset \mathcal{E}$  such that*

$$F(\mathbf{X}) := (P_{\mathcal{E}} \otimes I) \mathcal{P}(\mathbf{V}, \mathbf{X})|_{\mathcal{E} \otimes \mathcal{H}}, \quad \mathbf{X} \in \mathbf{B}_n(\mathcal{H}).$$