

CHOQUET ORDER AND HYPERRIGIDITY FOR FUNCTION SYSTEMS

Kenneth R. Davidson

University of Waterloo

Haifa, June 2017

joint work with [Matthew Kennedy](#)

$1 \in \mathcal{S} = \mathcal{S}^* \subset C(X)$ is a **function system**.

$K = \{\varphi \in \mathcal{S}^* : \varphi \geq 0, \varphi(1) = 1\}$ **state space**, compact, convex,
 and $x \in X \rightarrow \varepsilon_x \in K$, where $\varepsilon_x(f) = f(x)$ for $f \in \mathcal{S}$.

$1 \in \mathcal{S} = \mathcal{S}^* \subset C(X)$ is a **function system**.

$K = \{\varphi \in \mathcal{S}^* : \varphi \geq 0, \varphi(1) = 1\}$ **state space**, compact, convex,
 and $x \in X \rightarrow \varepsilon_x \in K$, where $\varepsilon_x(f) = f(x)$ for $f \in \mathcal{S}$.

THEOREM (KADISON 1951)

$\mathcal{S} \xrightarrow{\text{iso}} A(K) \subset C(K)$ isometric isomorphism to affine functions.

$1 \in \mathcal{S} = \mathcal{S}^* \subset C(X)$ is a **function system**.

$K = \{\varphi \in \mathcal{S}^* : \varphi \geq 0, \varphi(1) = 1\}$ **state space**, compact, convex,
 and $x \in X \rightarrow \varepsilon_x \in K$, where $\varepsilon_x(f) = f(x)$ for $f \in \mathcal{S}$.

THEOREM (KADISON 1951)

$\mathcal{S} \xrightarrow{\text{iso}} A(K) \subset C(K)$ isometric isomorphism to affine functions.

$\partial\mathcal{S} := \partial K$ extreme points is **Choquet boundary** of \mathcal{S} .

$f \in \mathcal{S}$ affine on K , so $\mathcal{S} \rightarrow C(\overline{\partial K})$ completely isometric.

$\overline{\partial K}$ is the **Shilov boundary** of \mathcal{S} .

By Hahn-Banach and Riesz Representation Theorems,
for $\varphi \in K$ there exists $\mu \in M_+(\overline{\partial K})$ **representing measure**:

$$\varphi(f) = \int f d\mu \quad \text{for } f \in \mathcal{S}.$$

By Hahn-Banach and Riesz Representation Theorems,
 for $\varphi \in K$ there exists $\mu \in M_+(\overline{\partial K})$ **representing measure**:

$$\varphi(f) = \int f d\mu \quad \text{for } f \in \mathcal{S}.$$

Choquet theory yields $\mu \in M_+(\partial K)$.

- important in applications
- nonmetrizable case: ∂K may not be Borel;
 so need technical definition of support.

By Hahn-Banach and Riesz Representation Theorems,
 for $\varphi \in K$ there exists $\mu \in M_+(\overline{\partial K})$ **representing measure**:

$$\varphi(f) = \int f d\mu \quad \text{for } f \in \mathcal{S}.$$

Choquet theory yields $\mu \in M_+(\partial K)$.

- important in applications
- nonmetrizable case: ∂K may not be Borel;
 so need technical definition of support.

DEFINITION

Choquet order: $\mu \prec_c \nu$ in $M_+(K)$ if $\int f d\mu \leq \int f d\nu$ for f convex.

This implies that $\int f d\mu = \int f d\nu$ for $f \in \mathcal{S}$, so represent same φ .

THEOREM (CHOQUET, MOKOBODSKI)

K metrizable.

$\mu \in M_+(K)$ is maximal in $\prec_c \iff \text{supp } \mu \subset \partial K$.

THEOREM (CHOQUET, MOKOBODSKI)

K metrizable.

$\mu \in M_+(K)$ is maximal in $\prec_c \iff \text{supp } \mu \subset \partial K$.

THEOREM (BISHOP-DE LEUW)

K arbitrary.

μ is maximal in $\prec_c \implies \mu(A) = 0$ if A Baire s.t. $A \cap \partial K = \emptyset$.

THEOREM (CHOQUET, MOKOBODSKI)

K metrizable.

$\mu \in M_+(K)$ is maximal in $\prec_c \iff \text{supp } \mu \subset \partial K$.

THEOREM (BISHOP-DE LEUW)

K arbitrary.

μ is maximal in $\prec_c \implies \mu(A) = 0$ if A Baire s.t. $A \cap \partial K = \emptyset$.

Mokobodski: this does not characterize maximality.

However, if ∂K is closed, then μ is maximal $\iff \text{supp } \mu \subset \partial K$.

Classical result:

THEOREM (KOROVKIN)

If $\Phi_n : C[a, b] \rightarrow C[a, b]$ positive maps s.t.
$$\lim_{n \rightarrow \infty} \Phi_n(f) = f \quad \text{for } f \in \{1, x, x^2\},$$

then
$$\lim_{n \rightarrow \infty} \Phi_n(f) = f \quad \text{for all } f \in C[a, b].$$

Classical result:

THEOREM (KOROVKIN)

If $\Phi_n : C[a, b] \rightarrow C[a, b]$ positive maps s.t.

$$\lim_{n \rightarrow \infty} \Phi_n(f) = f \quad \text{for } f \in \{1, x, x^2\},$$
 then
$$\lim_{n \rightarrow \infty} \Phi_n(f) = f \quad \text{for all } f \in C[a, b].$$

modern, significant improvement:

THEOREM (ARVESON)

If $\pi : C[a, b] \rightarrow \mathcal{B}(\mathcal{H})$ $*$ -repn., $\Phi_n : C[a, b] \rightarrow \mathcal{B}(\mathcal{H})$ (completely) positive maps s.t.

$$\lim_{n \rightarrow \infty} \Phi_n(f) = \pi(f) \quad \text{for } f \in \{1, x, x^2\},$$

then
$$\lim_{n \rightarrow \infty} \Phi_n(f) = \pi(f) \quad \text{for all } f \in C[a, b].$$

DEFINITION

$1 \in F \subset C(X)$ is a **Korovkin set** if $\Phi_n : C(X) \rightarrow C(X)$ are positive,
 $\lim_{n \rightarrow \infty} \Phi_n(f) = f$ for $f \in F \implies \lim_{n \rightarrow \infty} \Phi_n(f) = f$ for $f \in C(X)$.

DEFINITION

$1 \in F \subset C(X)$ is a **Korovkin set** if $\Phi_n : C(X) \rightarrow C(X)$ are positive,
 $\lim_{n \rightarrow \infty} \Phi_n(f) = f$ for $f \in F \implies \lim_{n \rightarrow \infty} \Phi_n(f) = f$ for $f \in C(X)$.

F is a **strong Korovkin set** if $\pi : C[a, b] \rightarrow \mathcal{B}(\mathcal{H})$ $*$ -repn.,
 $\Phi_n : C(X) \rightarrow \mathcal{B}(\mathcal{H})$ (completely) positive, then
 $\lim_{n \rightarrow \infty} \Phi_n(f) = \pi(f)$ for $f \in F \implies \lim_{n \rightarrow \infty} \Phi_n(f) = \pi(f)$ for $f \in C(X)$.

DEFINITION

$1 \in F \subset C(X)$ is a **Korovkin set** if $\Phi_n : C(X) \rightarrow C(X)$ are positive,
 $\lim_{n \rightarrow \infty} \Phi_n(f) = f$ for $f \in F \implies \lim_{n \rightarrow \infty} \Phi_n(f) = f$ for $f \in C(X)$.

F is a **strong Korovkin set** if $\pi : C[a, b] \rightarrow \mathcal{B}(\mathcal{H})$ *-repn.,
 $\Phi_n : C(X) \rightarrow \mathcal{B}(\mathcal{H})$ (completely) positive, then
 $\lim_{n \rightarrow \infty} \Phi_n(f) = \pi(f)$ for $f \in F \implies \lim_{n \rightarrow \infty} \Phi_n(f) = \pi(f)$ for $f \in C(X)$.

THEOREM (ŠAŠKIN)

X compact metric. $1 \in F \subset C(X)$. $\mathcal{S} = \overline{\text{span}}\{F \cup F^*\}$.
 Then F is a Korovkin set $\iff \partial\mathcal{S} = X$.

DEFINITION

$1 \in F \subset C(X)$ is a **Korovkin set** if $\Phi_n : C(X) \rightarrow C(X)$ are positive,
 $\lim_{n \rightarrow \infty} \Phi_n(f) = f$ for $f \in F \implies \lim_{n \rightarrow \infty} \Phi_n(f) = f$ for $f \in C(X)$.

F is a **strong Korovkin set** if $\pi : C[a, b] \rightarrow \mathcal{B}(\mathcal{H})$ *-repn.,
 $\Phi_n : C(X) \rightarrow \mathcal{B}(\mathcal{H})$ (completely) positive, then
 $\lim_{n \rightarrow \infty} \Phi_n(f) = \pi(f)$ for $f \in F \implies \lim_{n \rightarrow \infty} \Phi_n(f) = \pi(f)$ for $f \in C(X)$.

THEOREM (ŠAŠKIN)

X compact metric. $1 \in F \subset C(X)$. $\mathcal{S} = \overline{\text{span}}\{F \cup F^*\}$.
 Then F is a Korovkin set $\iff \partial\mathcal{S} = X$.

QUESTION (ARVESON)

Characterize strong Korovkin sets.

DEFINITION

$1 \in \mathcal{S} = \mathcal{S}^* \subset \mathfrak{A} = C^*(\mathcal{S})$ is **hyperrigid** if whenever

$\pi : \mathfrak{A} \rightarrow \mathcal{B}(\mathcal{H})$ $*$ -repn, and $\Phi_n : \mathfrak{A} \rightarrow \mathcal{B}(\mathcal{H})$ c.p.

$\lim_{n \rightarrow \infty} \Phi_n(s) = \pi(s)$ for $s \in \mathcal{S} \implies \lim_{n \rightarrow \infty} \Phi_n(a) = \pi(a)$ for $a \in \mathfrak{A}$.

DEFINITION

$1 \in \mathcal{S} = \mathcal{S}^* \subset \mathfrak{A} = C^*(\mathcal{S})$ is **hypperrigid** if whenever

$\pi : \mathfrak{A} \rightarrow \mathcal{B}(\mathcal{H})$ $*$ -repn, and $\Phi_n : \mathfrak{A} \rightarrow \mathcal{B}(\mathcal{H})$ c.p.

$\lim_{n \rightarrow \infty} \Phi_n(s) = \pi(s)$ for $s \in \mathcal{S} \implies \lim_{n \rightarrow \infty} \Phi_n(a) = \pi(a)$ for $a \in \mathfrak{A}$.

DEFINITION

$1 \in \mathcal{S} = \mathcal{S}^* \subset \mathfrak{A} = C^*(\mathcal{S})$. $\pi : \mathfrak{A} \rightarrow \mathcal{B}(\mathcal{H})$ $*$ -repn.

$\pi|_{\mathcal{S}}$ has **unique extension property (u.e.p.)**

if π is the unique u.c.p. extension to \mathfrak{A} .

DEFINITION

$1 \in \mathcal{S} = \mathcal{S}^* \subset \mathfrak{A} = C^*(\mathcal{S})$ is **hypermrigid** if whenever

$\pi : \mathfrak{A} \rightarrow \mathcal{B}(\mathcal{H})$ $*$ -repn, and $\Phi_n : \mathfrak{A} \rightarrow \mathcal{B}(\mathcal{H})$ c.p.

$\lim_{n \rightarrow \infty} \Phi_n(s) = \pi(s)$ for $s \in \mathcal{S} \implies \lim_{n \rightarrow \infty} \Phi_n(a) = \pi(a)$ for $a \in \mathfrak{A}$.

DEFINITION

$1 \in \mathcal{S} = \mathcal{S}^* \subset \mathfrak{A} = C^*(\mathcal{S})$. $\pi : \mathfrak{A} \rightarrow \mathcal{B}(\mathcal{H})$ $*$ -repn.

$\pi|_{\mathcal{S}}$ has **unique extension property (u.e.p.)**

if π is the unique u.c.p. extension to \mathfrak{A} .

THEOREM (ARVESON)

$1 \in \mathcal{S} = \mathcal{S}^* \subset \mathfrak{A} = C^*(\mathcal{S})$. Then

\mathcal{S} is hypermrigid $\iff \pi|_{\mathcal{S}}$ has u.e.p. $\forall \pi$ $*$ -repn.

DEFINITION

π *-repn. of \mathfrak{A} is a **boundary representation for \mathcal{S}** if
 π is irreducible and $\pi|_{\mathcal{S}}$ has u.e.p.

DEFINITION

π $*$ -repr. of \mathfrak{A} is a **boundary representation for \mathcal{S}** if
 π is irreducible and $\pi|_{\mathcal{S}}$ has u.e.p.

CONJECTURE (ARVESON)

\mathcal{S} is hyperrigid \iff every *irreducible* $*$ -repr. is a boundary repr.

DEFINITION

π $*$ -repr. of \mathfrak{A} is a **boundary representation for \mathcal{S}** if π is irreducible and $\pi|_{\mathcal{S}}$ has u.e.p.

CONJECTURE (ARVESON)

\mathcal{S} is hyperrigid \iff every *irreducible* $*$ -repr. is a boundary repr.

REMARK

For $1 \in \mathcal{S} = \mathcal{S}^* \subset C(X)$, this asks if $\partial\mathcal{S} = X$,
 is \mathcal{S} a strong Korovkin set in $C(\partial\mathcal{S})$?

DEFINITION

Dilation order: $\mu \prec_d \nu \in M_+(K)$ if there exist ***-repns.**

$$\pi : C(K) \rightarrow \mathcal{B}(\mathcal{H}), \quad \xi \in \mathcal{H}, \quad \langle \pi(f)\xi, \xi \rangle = \int f d\mu \quad \forall f \in C(K)$$

$$\sigma : C(K) \rightarrow \mathcal{B}(\mathcal{K}), \quad \eta \in \mathcal{K}, \quad \langle \sigma(f)\eta, \eta \rangle = \int f d\nu \quad \forall f \in C(K)$$

and **isometry** $J : \mathcal{H} \rightarrow \mathcal{K}$ s.t. $J\xi = \eta$ and $J^*\sigma(f)J = \pi(f) \quad \forall f \in \mathcal{S}$.

DEFINITION

Dilation order: $\mu \prec_d \nu \in M_+(K)$ if there exist ***-repns.**

$$\pi : C(K) \rightarrow \mathcal{B}(\mathcal{H}), \quad \xi \in \mathcal{H}, \quad \langle \pi(f)\xi, \xi \rangle = \int f d\mu \quad \forall f \in C(K)$$

$$\sigma : C(K) \rightarrow \mathcal{B}(\mathcal{K}), \quad \eta \in \mathcal{K}, \quad \langle \sigma(f)\eta, \eta \rangle = \int f d\nu \quad \forall f \in C(K)$$

and **isometry** $J : \mathcal{H} \rightarrow \mathcal{K}$ s.t. $J\xi = \eta$ and $J^*\sigma(f)J = \pi(f) \quad \forall f \in \mathcal{S}$.

THEOREM 1

Dilation order is the same as Choquet order.

DEFINITION

Dilation order: $\mu \prec_d \nu \in M_+(K)$ if there exist ***-repns.**

$$\pi : C(K) \rightarrow \mathcal{B}(\mathcal{H}), \quad \xi \in \mathcal{H}, \quad \langle \pi(f)\xi, \xi \rangle = \int f d\mu \quad \forall f \in C(K)$$

$$\sigma : C(K) \rightarrow \mathcal{B}(\mathcal{K}), \quad \eta \in \mathcal{K}, \quad \langle \sigma(f)\eta, \eta \rangle = \int f d\nu \quad \forall f \in C(K)$$

and **isometry** $J : \mathcal{H} \rightarrow \mathcal{K}$ s.t. $J\xi = \eta$ and $J^*\sigma(f)J = \pi(f) \quad \forall f \in \mathcal{S}$.

THEOREM 1

Dilation order is the same as Choquet order.

COROLLARY

$\mu \prec_c \nu \iff \exists \Phi : C(K) \rightarrow L^\infty(\mu)$ positive s.t.

- ① $\Phi(f) = f$ for all $f \in A(K)$, and
- ② $\int \Phi(f) d\mu = \int f d\nu$ for all $f \in C(K)$.

$\pi_\mu : C(K) \rightarrow B(L^2(\mu))$ by $\pi(f) = M_f$.

THEOREM 2

π_μ has u.e.p. $\iff \mu$ is maximal in \prec_d .

$\pi_\mu : C(K) \rightarrow B(L^2(\mu))$ by $\pi(f) = M_f$.

THEOREM 2

π_μ has u.e.p. $\iff \mu$ is maximal in \prec_d .

COROLLARY (HYPERRIGIDITY FOR FUNCTION SYSTEMS)

If ∂S is closed, then S is hyperrigid in $C(\partial S)$.

$\pi_\mu : C(K) \rightarrow \mathcal{B}(L^2(\mu))$ by $\pi(f) = M_f$.

THEOREM 2

π_μ has u.e.p. $\iff \mu$ is maximal in \prec_d .

COROLLARY (HYPERRIGIDITY FOR FUNCTION SYSTEMS)

If $\partial\mathcal{S}$ is closed, then \mathcal{S} is hyperrigid in $C(\partial\mathcal{S})$.

COROLLARY

If X is metrizable, $1 = \mathcal{S} \subset C(X)$, $\pi : C(X) \rightarrow \mathcal{B}(\mathcal{H})$ $*$ -repn.
 Then π has u.e.p. $\iff \pi$ is supported on $\partial\mathcal{S}$.

Application to approximation theory

The following does not require metrizability, so it generalizes Šaškin's Theorem even in the classical situation.

COROLLARY

$1 \in \mathcal{S} = \overline{\text{span}}\{F \cup F^*\} \subset C(X)$.

TFAE

- ① $\partial\mathcal{S} = X$
- ② F is a Korovkin set.
- ③ F is a strong Korovkin set.

Application to classical Choquet theory

THEOREM (CARTIER)

If K is metrizable, $\mu \prec_c \nu$, then $\exists \lambda : K \rightarrow M_{+,1}(K)$ s.t.

- ① $x \rightarrow \lambda_x(f)$ is Borel $\forall f \in C(K)$,
- ② $\lambda_x(f) = f(x) \quad \forall f \in A(K)$, and
- ③ $\int f d\nu = \int \lambda_x(f) d\mu \quad \forall f \in C(K)$.

Application to classical Choquet theory

THEOREM (CARTIER)

If K is metrizable, $\mu \prec_c \nu$, then $\exists \lambda : K \rightarrow M_{+,1}(K)$ s.t.

- ① $x \rightarrow \lambda_x(f)$ is Borel $\forall f \in C(K)$,
- ② $\lambda_x(f) = f(x) \quad \forall f \in A(K)$, and
- ③ $\int f d\nu = \int \lambda_x(f) d\mu \quad \forall f \in C(K)$.

THEOREM 3

K compact convex, $\mu \prec_c \nu$, then $\exists \lambda : K \rightarrow M_{+,1}(K)$ s.t.

- ① $x \rightarrow \lambda_x(f)$ is Borel $\forall f \in C(K)$,
- ② $\lambda_x(f) = f(x)$ a.e. $(\mu) \quad \forall f \in A(K)$, and
- ③ $\int f d\nu = \int \lambda_x(f) d\mu \quad \forall f \in C(K)$.

Thank you.
The end.